

## FORECAST COMBINATIONS

ALLAN TIMMERMANN

UCSD

### Contents

Abstract	136
Keywords	136
1. Introduction	137
2. The forecast combination problem	140
2.1. Specification of loss function	141
2.2. Construction of a super model – pooling information	143
2.3. Linear forecast combinations under MSE loss	144
2.3.1. Diversification gains	145
2.3.2. Effect of bias in individual forecasts	148
2.4. Optimality of equal weights – general case	148
2.5. Optimal combinations under asymmetric loss	150
2.6. Combining as a hedge against non-stationarities	154
3. Estimation	156
3.1. To combine or not to combine	156
3.2. Least squares estimators of the weights	158
3.3. Relative performance weights	159
3.4. Moment estimators	160
3.5. Nonparametric combination schemes	160
3.6. Pooling, clustering and trimming	162
4. Time-varying and nonlinear combination methods	165
4.1. Time-varying weights	165
4.2. Nonlinear combination schemes	169
5. Shrinkage methods	170
5.1. Shrinkage and factor structure	172
5.2. Constraints on combination weights	174
6. Combination of interval and probability distribution forecasts	176
6.1. The combination decision	176
6.2. Combinations of probability density forecasts	177
6.3. Bayesian methods	178

6.3.1. Bayesian model averaging	179
6.4. Combinations of quantile forecasts	179
7. Empirical evidence	181
7.1. Simple combination schemes are hard to beat	181
7.2. Choosing the single forecast with the best track record is often a bad idea	182
7.3. Trimming of the worst models often improves performance	183
7.4. Shrinkage often improves performance	184
7.5. Limited time-variation in the combination weights may be helpful	185
7.6. Empirical application	186
8. Conclusion	193
Acknowledgements	193
References	194

## Abstract

Forecast combinations have frequently been found in empirical studies to produce better forecasts on average than methods based on the *ex ante* best individual forecasting model. Moreover, simple combinations that ignore correlations between forecast errors often dominate more refined combination schemes aimed at estimating the theoretically optimal combination weights. In this chapter we analyze theoretically the factors that determine the advantages from combining forecasts (for example, the degree of correlation between forecast errors and the relative size of the individual models' forecast error variances). Although the reasons for the success of simple combination schemes are poorly understood, we discuss several possibilities related to model misspecification, instability (non-stationarities) and estimation error in situations where the number of models is large relative to the available sample size. We discuss the role of combinations under asymmetric loss and consider combinations of point, interval and probability forecasts.

## Keywords

forecast combinations, pooling and trimming, shrinkage methods, model misspecification, diversification gains

*JEL classification:* C53, C22

## 1. Introduction

Multiple forecasts of the same variable are often available to decision makers. This could reflect differences in forecasters' subjective judgements due to heterogeneity in their information sets in the presence of private information or due to differences in modelling approaches. In the latter case, two forecasters may well arrive at very different views depending on the maintained assumptions underlying their forecasting models, e.g., constant versus time-varying parameters, linear versus nonlinear forecasting models, etc.

Faced with multiple forecasts of the same variable, an issue that immediately arises is how best to exploit information in the individual forecasts. In particular, should a single dominant forecast be identified or should a combination of the underlying forecasts be used to produce a pooled summary measure? From a theoretical perspective, unless one can identify *ex ante* a particular forecasting model that generates smaller forecast errors than its competitors (and whose forecast errors cannot be hedged by other models' forecast errors), forecast combinations offer diversification gains that make it attractive to combine individual forecasts rather than relying on forecasts from a single model. Even if the best model could be identified at each point in time, combination may still be an attractive strategy due to diversification gains, although its success will depend on how well the combination weights can be determined.

Forecast combinations have been used successfully in empirical work in such diverse areas as forecasting Gross National Product, currency market volatility, inflation, money supply, stock prices, meteorological data, city populations, outcomes of football games, wilderness area use, check volume and political risks, cf. [Clemen \(1989\)](#). Summarizing the simulation and empirical evidence in the literature on forecast combinations, [Clemen \(1989, p. 559\)](#) writes "The results have been virtually unanimous: combining multiple forecasts leads to increased forecast accuracy . . . in many cases one can make dramatic performance improvements by simply averaging the forecasts." More recently, [Makridakis and Hibon \(2000\)](#) conducted the so-called M3-competition which involved forecasting 3003 time series and concluded (p. 458) "The accuracy of the combination of various methods outperforms, on average, the specific methods being combined and does well in comparison with other methods." Similarly, [Stock and Watson \(2001, 2004\)](#) undertook an extensive study across numerous economic and financial variables using linear and nonlinear forecasting models and found that, on average, pooled forecasts outperform predictions from the single best model, thus confirming Clemen's conclusion. Their analysis has been extended to a large European data set by [Marcellino \(2004\)](#) with essentially the same conclusions.

A simple portfolio diversification argument motivates the idea of combining forecasts, cf. [Bates and Granger \(1969\)](#). Its premise is that, perhaps due to presence of private information, the information set underlying the individual forecasts is often unobserved to the forecast user. In this situation it is not feasible to pool the underlying information sets and construct a 'super' model that nests each of the underlying forecasting models. For example, suppose that we are interested in forecasting some variable,  $y$ ,

and that two predictions,  $\hat{y}_1$  and  $\hat{y}_2$  of its conditional mean are available. Let the first forecast be based on the variables  $x_1, x_2$ , i.e.,  $\hat{y}_1 = g_1(x_1, x_2)$ , while the second forecast is based on the variables  $x_3, x_4$ , i.e.,  $\hat{y}_2 = g_2(x_3, x_4)$ . Further, suppose that all variables enter with non-zero weights in the forecasts and that the  $x$ -variables are imperfectly correlated. If  $\{x_1, x_2, x_3, x_4\}$  were observable, it would be natural to construct a forecasting model based on all four variables,  $\hat{y}_3 = g_3(x_1, x_2, x_3, x_4)$ . On the other hand, if only the forecasts,  $\hat{y}_1$  and  $\hat{y}_2$  are observed by the forecast user (while the underlying variables are unobserved) then the only option is to combine these forecasts, i.e. to elicit a model of the type  $\hat{y} = g_c(\hat{y}_1, \hat{y}_2)$ . More generally, the forecast user's information set,  $\mathcal{F}$ , may comprise  $n$  individual forecasts,  $\mathcal{F} = \{\hat{y}_1, \dots, \hat{y}_n\}$ , where  $\mathcal{F}$  is often not the union of the information sets underlying the individual forecasts,  $\bigcup_{i=1}^n \mathcal{F}_i$ , but a much smaller subset. Of course, the higher the degree of overlap in the information sets used to produce the underlying forecasts, the less useful a combination of forecasts is likely to be, cf. Clemen (1987).

It is difficult to fully appreciate the strength of the diversification or hedging argument underlying forecast combination. Suppose the aim is to minimize some loss function belonging to a family of convex loss functions,  $\mathcal{L}$ , and that some forecast,  $\hat{y}_1$ , stochastically dominates another forecast,  $\hat{y}_2$ , in the sense that expected losses for all loss functions in  $\mathcal{L}$  are lower under  $\hat{y}_1$  than under  $\hat{y}_2$ . While this means that it is not rational for a decision maker to choose  $\hat{y}_2$  over  $\hat{y}_1$  in isolation, it is easy to construct examples where some combination of  $\hat{y}_1$  and  $\hat{y}_2$  generates a smaller expected loss than that produced using  $\hat{y}_1$  alone.

A second reason for using forecast combinations referred to by, inter alia, Figlewski and Ulrich (1983), Kang (1986), Diebold and Pauly (1987), Makridakis (1989), Sessions and Chattererjee (1989), Winkler (1989), Hendry and Clements (2002) and Aiolfi and Timmermann (2006) and also thought of by Bates and Granger (1969), is that individual forecasts may be very differently affected by structural breaks caused, for example, by institutional change or technological developments. Some models may adapt quickly and will only temporarily be affected by structural breaks, while others have parameters that only adjust very slowly to new post-break data. The more data that is available after the most recent break, the better one might expect stable, slowly adapting models to perform relative to fast adapting ones as the parameters of the former are more precisely estimated. Conversely, if the data window since the most recent break is short, the faster adapting models can be expected to produce the best forecasting performance. Since it is typically difficult to detect structural breaks in 'real time', it is plausible that on average, i.e., across periods with varying degrees of stability, combinations of forecasts from models with different degrees of adaptability will outperform forecasts from individual models. This intuition is confirmed in Pesaran and Timmermann (2005).

A third and related reason for forecast combination is that individual forecasting models may be subject to misspecification bias of unknown form, a point stressed particularly by Clemen (1989), Makridakis (1989), Diebold and Lopez (1996) and Stock and Watson (2001, 2004). Even in a stationary world, the true data generating process is likely to be more complex and of a much higher dimension than assumed by the

most flexible and general model entertained by a forecaster. Viewing forecasting models as local approximations, it is implausible that the same model dominates all others at all points in time. Rather, the best model may change over time in ways that can be difficult to track on the basis of past forecasting performance. Combining forecasts across different models can be viewed as a way to make the forecast more robust against such misspecification biases and measurement errors in the data sets underlying the individual forecasts. Notice again the similarity to the classical portfolio diversification argument for risk reduction: Here the portfolio is the combination of forecasts and the source of risk reflects incomplete information about the target variable and model misspecification possibly due to non-stationarities in the underlying data generating process.

A fourth argument for combination of forecasts is that the underlying forecasts may be based on different loss functions. This argument holds even if the forecasters observe the same information set. Suppose, for example, that forecaster A strongly dislikes large negative forecast errors while forecaster B strongly dislikes large positive forecast errors. In this case, forecaster A is likely to under-predict the variable of interest (so the forecast error distribution is centered on a positive value), while forecaster B will over-predict it. If the bias is constant over time, there is no need to average across different forecasts since including a constant in the combination equation will pick up any unwanted bias. Suppose, however, that the optimal amount of bias is proportional to the conditional variance of the variable, as in [Christoffersen and Diebold \(1997\)](#) and [Zellner \(1986\)](#). Provided that the two forecasters adopt a similar volatility model (which is not implausible since they are assumed to share the same information set), a forecast user with a more symmetric loss function than was used to construct the underlying forecasts could find a combination of the two forecasts better than the individual ones.

Numerous arguments against using forecast combinations can also be advanced. Estimation errors that contaminate the combination weights are known to be a serious problem for many combination techniques especially when the sample size is small relative to the number of forecasts, cf. [Diebold and Pauly \(1990\)](#), [Elliott \(2004\)](#) and [Yang \(2004\)](#). Although non-stationarities in the underlying data generating process can be an argument for using combinations, it can also lead to instabilities in the combination weights and lead to difficulties in deriving a set of combination weights that performs well, cf. [Clemen and Winkler \(1986\)](#), [Diebold and Pauly \(1987\)](#), [Figlewski and Urich \(1983\)](#), [Kang \(1986\)](#) and [Palm and Zellner \(1992\)](#). In situations where the information sets underlying the individual forecasts are unobserved, most would agree that forecast combinations can add value. However, when the full set of predictor variables used to construct different forecasts is observed by the forecast user, the use of a combination strategy instead of attempting to identify a single best “super” model can be challenged, cf. [Chong and Hendry \(1986\)](#) and [Diebold \(1989\)](#).

It is no coincidence that these arguments against forecast combinations seem familiar. In fact, there are many similarities between the forecast combination problem and the standard problem of constructing a single econometric specification. In both cases a subset of predictors (or individual forecasts) has to be selected from a larger set of potential

forecasting variables and the choice of functional form mapping this information into the forecast as well as the choice of estimation method have to be determined. There are clearly important differences as well. First, it may be reasonable to assume that the individual forecasts are unbiased in which case the combined forecast will also be unbiased provided that the combination weights are constrained to sum to unity and an intercept is omitted. Provided that the unbiasedness assumption holds for each forecast, imposing such parameter constraints can lead to efficiency gains. One would almost never want to impose this type of constraint on the coefficients of a standard regression model since predictor variables can differ significantly in their units, interpretation and scaling. Secondly, if the individual forecasts are generated by quantitative models whose parameters are estimated recursively there is a potential generated regressor problem which could bias estimates of the combination weights. In part this explains why using simple averages based on equal weights provides a natural benchmark. Finally, the forecasts that are being combined need not be point forecasts but could take the form of interval or density forecasts.

As a testimony to its important role in the forecasting literature, many high-quality surveys of forecast combinations have already appeared, cf. [Clemen \(1989\)](#), [Diebold and Lopez \(1996\)](#) and [Newbold and Harvey \(2001\)](#). This survey differs from earlier ones in many important ways, however. First, we put more emphasis on the theory underlying forecast combinations, particularly in regard to the diversification argument which is common also in portfolio analysis. Second, we deal in more depth with recent topics – some of which were emphasized as important areas of future research by [Diebold and Lopez \(1996\)](#) – such as combination of probability forecasts, time-varying combination weights, combination under asymmetric loss and shrinkage.

The chapter is organized as follows. We first develop the theory underlying the general forecast combination problem in Section 2. The following section discusses estimation methods for the linear forecast combination problem. Section 4 considers nonlinear combination schemes and combinations with time-varying weights. Section 5 discusses shrinkage combinations while Section 6 covers combinations of interval or density forecasts. Section 7 extracts main conclusions from the empirical literature and Section 8 concludes.

## 2. The forecast combination problem

Consider the problem of forecasting at time  $t$  the future value of some target variable,  $y$ , after  $h$  periods, whose realization is denoted  $y_{t+h}$ . Since no major new insights arise from the case where  $y$  is multivariate, to simplify the exposition we shall assume that  $y_{t+h} \in \mathbb{R}$ . We shall refer to  $t$  as the time of the forecast and  $h$  as the forecast horizon. The information set at time  $t$  will be denoted by  $\mathcal{F}_t$  and we assume that  $\mathcal{F}_t$  comprises an  $N$  – vector of forecasts  $\hat{\mathbf{y}}_{t+h,t} = (\hat{y}_{t+h,t,1}, \hat{y}_{t+h,t,2}, \dots, \hat{y}_{t+h,t,N})'$  in addition to the histories of these forecasts up to time  $t$  and the history of the realizations of the

target variable, i.e.  $\mathcal{F}_t = \{\hat{\mathbf{y}}_{h+1,1}, \hat{\mathbf{y}}_{t+h,t}, y_1, \dots, y_t\}$ . A set of additional information variables,  $\mathbf{x}_t$ , can easily be included in the problem.

The general forecast combination problem seeks an aggregator that reduces the information in a potentially high-dimensional vector of forecasts,  $\hat{\mathbf{y}}_{t+h,t} \in \mathbb{R}^N$ , to a lower dimensional summary measure,  $C(\hat{\mathbf{y}}_{t+h,t}; \boldsymbol{\omega}_c) \in \mathbb{R}^c \subset \mathbb{R}^N$ , where  $\boldsymbol{\omega}_c$  are the parameters associated with the combination. If only a point forecast is of interest, then a one-dimensional aggregator will suffice. For example, a decision maker interested in using forecasts to determine how much to invest in a risky asset may want to use not only information on either the mode, median or mean forecast, but also to consider the degree of dispersion across individual forecasts as a way to measure the uncertainty or ‘disagreement’ surrounding the forecasts. How low-dimensional the combined forecast should be is not always obvious. Outside the MSE framework, it is not trivially true that a scalar aggregator that summarizes all relevant information can always be found.

Forecasts do not intrinsically have direct value to decision makers. Rather, they become valuable only to the extent that they can be used to improve decision makers’ actions, which in turn affect their loss or utility. Point forecasts generally provide insufficient information for a decision maker or forecast user who, for example, may be interested in the degree of uncertainty surrounding the forecast. Nevertheless, the vast majority of studies on forecast combinations has dealt with point forecasts so we initially focus on this case. We let  $\hat{y}_{t+h,t}^c = C(\hat{\mathbf{y}}_{t+h,t}; \boldsymbol{\omega}_{t+h,t})$  be the combined point forecast as a function of the underlying forecasts  $\hat{\mathbf{y}}_{t+h,t}$  and the parameters of the combination,  $\boldsymbol{\omega}_{t+h,t} \in \mathcal{W}_t$ , where  $\mathcal{W}_t$  is often assumed to be a compact subset of  $\mathbb{R}^N$  and  $\boldsymbol{\omega}_{t+h,t}$  can be time-varying but is adapted to  $\mathcal{F}_t$ . For example, equal weights would give  $g(\hat{\mathbf{y}}_{t+h,t}; \boldsymbol{\omega}_{t+h,t}) = (1/N) \sum_{j=1}^N \hat{y}_{t+h,t,j}$ . Our choice of notation reflects that we will mostly be thinking of  $\boldsymbol{\omega}_{t+h,t}$  as combination weights, although the parameters need not always have this interpretation.

### 2.1. Specification of loss function

To simplify matters we follow standard practice and assume that the loss function only depends on the forecast error from the combination,  $e_{t+h,t}^c = y_{t+h} - g(\hat{\mathbf{y}}_{t+h,t}; \boldsymbol{\omega}_{t+h,t})$ , i.e.  $L = L(e_{t+h,t})$ . The vast majority of work on forecast combinations assumes this type of loss, in part because point forecasts are far more common than distribution forecasts and in part because the decision problem underlying the forecast situation is not worked out in detail. However, it should also be acknowledged that this loss function embodies a set of restrictive assumptions on the decision problem, cf. Granger and Machina (2006) and Elliott and Timmermann (2004). In Section 6 we cover the more general case that combines interval or distribution forecasts.

The parameters of the optimal combination,  $\boldsymbol{\omega}_{t+h,t}^* \in \mathcal{W}_t$ , solve the problem

$$\boldsymbol{\omega}_{t+h,t}^* = \arg \min_{\boldsymbol{\omega}_{t+h,t} \in \mathcal{W}_t} \mathbb{E}[L(e_{t+h,t}^c(\boldsymbol{\omega}_{t+h,t})) | \mathcal{F}_t]. \quad (1)$$

Here the expectation is taken over the conditional distribution of  $e_{t+h,t}$  given  $\mathcal{F}_t$ . Clearly optimality is established within the assumed family  $\hat{y}_{t+h,t}^c = C(\hat{\mathbf{y}}_{t+h,t}; \boldsymbol{\omega}_{t+h,t})$ . Elliott

and Timmermann (2004) show that, subject to a set of weak technical assumptions on the loss and distribution functions, the combination weights can be found as the solution to the following Taylor series expansion around  $\mu_{e_{t+h,t}} = E[e_{t+h,t} | \mathcal{F}_t]$ :

$$\omega_{t+h,t}^* = \arg \min_{\omega_{t+h,t} \in \mathcal{W}_t} \left\{ L(\mu_{e_{t+h,t}}) + \frac{1}{2} L''_{\mu_e} E[(e_{t+h,t} - \mu_{e_{t+h,t}})^2 | \mathcal{F}_t] + \sum_{m=3}^{\infty} L^m_{\mu_e} \sum_{i=0}^m \frac{1}{i!(m-i)!} E[e_{t+h,t}^{m-i} \mu_{e_{t+h,t}}^i | \mathcal{F}_t] \right\} \quad (2)$$

where  $L^k_{\mu_e} \equiv \partial^k L(e_{t+h,t}) / \partial^k e|_{e_{t+h,t} = \mu_{e_{t+h,t}}}$ . In general, the entire moment generating function of the forecast error distribution and all higher-order derivatives of the loss function will influence the optimal combination weights which therefore reflect both the shape of the loss function and the forecast error distribution.

The expansion in (2) suggests that the collection of individual forecasts  $\hat{y}_{t+h,t}$  is useful in as far as it can predict any of the conditional moments of the forecast error distribution of which a decision maker cares. Hence,  $\hat{y}_{t+h,t,i}$  gets a non-zero weight in the combination if for any moment,  $e_{t+h,t}^m$ , for which  $L^m_{\mu_e} \neq 0$ ,  $\partial E[e_{t+h,t}^m | \mathcal{F}_t] / \partial \hat{y}_{t+h,t,i} \neq 0$ . For example, if the vector of point forecasts can be used to predict the mean, variance, skew and kurtosis, but no other moments of the forecast error distribution, then the combined summary measure could be based on those summary measures of  $\hat{y}_{t+h,t}$  that predict the first through fourth moments.

Oftentimes it is simply assumed that the objective function underlying the combination problem is mean squared error (MSE) loss

$$L(y_{t+h}, \hat{y}_{t+h,t}) = \theta (y_{t+h} - \hat{y}_{t+h,t})^2, \quad \theta > 0. \quad (3)$$

For this case, the combined or consensus forecast seeks to choose a (possibly time-varying) mapping  $C(\hat{y}_{t+h,t}; \omega_{t+h,t})$  from the  $N$ -vector of individual forecasts  $\hat{y}_{t+h,t}$  to the real line,  $\mathcal{Y}_{t+h,t} \rightarrow \mathcal{R}$  that best approximates the conditional expectation,  $E[y_{t+h} | \hat{y}_{t+h,t}]$ .<sup>1</sup>

Two levels of aggregation are thus involved in the combination problem. The first step summarizes individual forecasters' private information to produce point forecasts  $\hat{y}_{t+h,t,i}$ . The only difference to the standard forecasting problem is that the 'input' variables are forecasts from other models or subjective forecasts. This may create a generated regressor problem that can bias the estimated combination weights, although this aspect is often ignored. It could in part explain why combinations based on estimated weights often do not perform well. The second step aggregates the vector of point forecasts  $\hat{y}_{t+h,t}$  to the consensus measure  $C(\hat{y}_{t+h,t}; \omega_{t+h,t})$ . Information is lost in both steps. Conversely, the second step is likely to lead to far simpler and more parsimonious forecasting models when compared to a forecast based on the full set of individual

<sup>1</sup> To see this, take expectations of (3) and differentiate with respect to  $C(\hat{y}_{t+h,t}; \omega_{t+h,t})$  to get  $C^*(\hat{y}_{t+h,t}; \omega_{t+h,t}) = E[y_{t+h} | \mathcal{F}_t]$ .



forecasts or a “super model” based on individual forecasters’ information variables. In general, we would expect information aggregation to increase the bias in the forecast but also to reduce the variance of the forecast error. To the extent possible, the combination should optimally trade off these two components. This is particularly clear under MSE loss, where the objective function equals the squared bias plus the forecast error variance,  $E[e_{t+h,t}^2] = E[e_{t+h,t}]^2 + \text{Var}(e_{t+h,t})$ .<sup>2</sup>

## 2.2. Construction of a super model – pooling information

Let  $\mathcal{F}_t^c = \bigcup_{i=1}^N \mathcal{F}_{it}$  be the union of the forecasters’ individual information sets, or the ‘super’ information set. If  $\mathcal{F}_t^c$  were observed, one possibility would be to model the conditional mean of  $y_{t+h}$  as a function of all these variables, i.e.

$$\hat{y}_{t+h,t} = C_s(\mathcal{F}_t^c; \boldsymbol{\theta}_{t+h,t,s}). \quad (4)$$

Individual forecasts,  $i$ , instead take the form  $\hat{y}_{t+h,t,i} = C_i(\mathcal{F}_{it}; \boldsymbol{\theta}_{t+h,t,i})$ .<sup>3</sup> If only the individual forecasts  $\hat{y}_{t+h,t,i}$  ( $i = 1, \dots, N$ ) are observed, whereas the underlying information sets  $\{\mathcal{F}_{it}\}$  are unobserved by the forecast user, the combined forecast would be restricted as follows:

$$\hat{y}_{t+h,t,i} = C_c(\hat{y}_{t+h,t,1}, \dots, \hat{y}_{t+h,t,N}; \boldsymbol{\theta}_{t+h,t,c}). \quad (5)$$

Normally it would be better to pool all information rather than first filter the information sets through the individual forecasting models. This introduces the usual efficiency loss through the two-stage estimation and also ignores correlations between the underlying information sources. There are several potential problems with pooling the information sets, however. One problem is – as already mentioned – that individual information sets may not be observable or too costly to combine. Diebold and Pauly (1990, p. 503) remark that “While pooling of forecasts is suboptimal relative to pooling of information sets, it must be recognized that in many forecasting situations, particularly in real time, pooling of information sets is either impossible or prohibitively costly.” Furthermore, in cases with many relevant input variables and complicated dynamic and nonlinear effects, constructing a “super model” using the pooled information set,  $\mathcal{F}_t^c$ , is not likely to provide good forecasts given the well-known problems associated with high-dimensional kernel regressions, nearest neighbor regressions, or other

<sup>2</sup> Clemen (1987) demonstrates that an important part of the aggregation of individual forecasts towards an aggregate forecast is an assessment of the dependence among the underlying models’ (‘experts’) forecasts and that a group forecast will generally be less informative than the set of individual forecasts. In fact, group forecasts only provide a sufficient statistic for collections of individual forecasts provided that both the experts and the decision maker agree in their assessments of the dependence among experts. This precludes differences in opinion about the correlation structure among decision makers. Taken to its extreme, this argument suggests that experts should not attempt to aggregate their observed information into a single forecast but should simply report their raw data to the decision maker.

<sup>3</sup> Notice that we use  $\boldsymbol{\omega}_{t+h,t}$  for the parameters involved in the combination of the forecasts,  $\hat{y}_{t+h,t}$ , while we use  $\boldsymbol{\theta}_{t+h,t}$  for the parameters relating the underlying information variables in  $\mathcal{F}_t$  to  $y_{t+h}$ .

nonparametric methods. Although individual forecasting models will be biased and may omit important variables, this bias can more than be compensated for by reductions in parameter estimation error in cases where the number of relevant predictor variables is much greater than  $N$ , the number of forecasts.<sup>4</sup>

### 2.3. Linear forecast combinations under MSE loss

While in general there is no closed-form solution to (1), one can get analytical results by imposing distributional restrictions or restrictions on the loss function. Unless the mapping,  $C$ , from  $\hat{\mathbf{y}}_{t+h,t}$  to  $y_{t+h}$  is modeled nonparametrically, optimality results for forecast combination must be established within families of parametric combination schemes of the form  $\hat{y}_{t+h,t}^c = C(\hat{\mathbf{y}}_{t+h,t}; \boldsymbol{\omega}_{t+h,t})$ . The general class of combination schemes in (1) comprises nonlinear as well as time-varying combination methods. We shall return to these but for now concentrate on the family of linear combinations,  $\mathcal{W}_t^l \subset \mathcal{W}_t$ , which are more commonly used.<sup>5</sup> To this end we choose weights,  $\boldsymbol{\omega}_{t+h,t} = (\omega_{t+h,t,1}, \dots, \omega_{t+h,t,N})'$  to produce a combined forecast of the form

$$\hat{y}_{t+h,t}^c = \boldsymbol{\omega}'_{t+h,t} \hat{\mathbf{y}}_{t+h,t}. \quad (6)$$

Under MSE loss, the combination weights are easy to characterize in population and only depend on the first two moments of the joint distribution of  $y_{t+h}$  and  $\hat{\mathbf{y}}_{t+h,t}$ ,

$$\begin{pmatrix} y_{t+h} \\ \hat{\mathbf{y}}_{t+h,t} \end{pmatrix} \sim \left( \begin{pmatrix} \mu_{y_{t+h,t}} \\ \boldsymbol{\mu}_{\hat{\mathbf{y}}_{t+h,t}} \end{pmatrix}, \begin{pmatrix} \sigma_{y_{t+h,t}}^2 & \boldsymbol{\sigma}'_{y\hat{\mathbf{y}}_{t+h,t}} \\ \boldsymbol{\sigma}_{y\hat{\mathbf{y}}_{t+h,t}} & \boldsymbol{\Sigma}_{\hat{\mathbf{y}}_{t+h,t}} \end{pmatrix} \right). \quad (7)$$

Minimizing  $E[e_{t+h,t}^2] = E[(y_{t+h} - \boldsymbol{\omega}'_{t+h,t} \hat{\mathbf{y}}_{t+h,t})^2]$ , we have

$$\begin{aligned} \boldsymbol{\omega}_{t+h,t}^* = \arg \min_{\boldsymbol{\omega}_{t+h,t} \in \mathcal{W}_t^l} & \left( (\mu_{y_{t+h,t}} - \boldsymbol{\omega}'_{t+h,t} \boldsymbol{\mu}_{\hat{\mathbf{y}}_{t+h,t}})^2 + \sigma_{y_{t+h,t}}^2 \right. \\ & \left. + \boldsymbol{\omega}'_{t+h,t} \boldsymbol{\Sigma}_{\hat{\mathbf{y}}_{t+h,t}} \boldsymbol{\omega}_{t+h,t} - 2\boldsymbol{\omega}'_{t+h,t} \boldsymbol{\sigma}_{y\hat{\mathbf{y}}_{t+h,t}} \right). \end{aligned}$$

This yields the first order condition

$$\begin{aligned} \frac{\partial E[e_{t+h,t}^2]}{\partial \boldsymbol{\omega}_{t+h,t}} &= -(\mu_{y_{t+h,t}} - \boldsymbol{\omega}'_{t+h,t} \boldsymbol{\mu}_{\hat{\mathbf{y}}_{t+h,t}}) \boldsymbol{\mu}_{\hat{\mathbf{y}}_{t+h,t}} + \boldsymbol{\Sigma}_{\hat{\mathbf{y}}_{t+h,t}} \boldsymbol{\omega}_{t+h,t} - \boldsymbol{\sigma}_{y\hat{\mathbf{y}}_{t+h,t}} \\ &= \mathbf{0}. \end{aligned}$$

Assuming that  $\boldsymbol{\Sigma}_{\hat{\mathbf{y}}_{t+h,t}}$  is invertible, this has the solution

$$\boldsymbol{\omega}_{t+h,t}^* = (\boldsymbol{\mu}_{\hat{\mathbf{y}}_{t+h,t}} \boldsymbol{\mu}'_{\hat{\mathbf{y}}_{t+h,t}} + \boldsymbol{\Sigma}_{\hat{\mathbf{y}}_{t+h,t}})^{-1} (\boldsymbol{\mu}_{\hat{\mathbf{y}}_{t+h,t}} \mu_{y_{t+h,t}} + \boldsymbol{\sigma}_{y\hat{\mathbf{y}}_{t+h,t}}). \quad (8)$$

<sup>4</sup> When the true forecasting model mapping  $\mathcal{F}_t^c$  to  $y_{t+h}$  is infinite-dimensional, the model that optimally balances bias and variance may depend on the sample size with a dimension that grows as the sample size increases.

<sup>5</sup> This, of course, does not rule out that the *estimated* weights vary over time as will be the case when the weights are updated recursively as more data becomes available.

This solution is optimal in population whenever  $y_{t+h}$  and  $\hat{y}_{t+h,t}$  are joint Gaussian since in this case the conditional expectation  $E[y_{t+h}|\hat{y}_{t+h,t}]$  will be linear in  $\hat{y}_{t+h,t}$ . For the moment we ignore time-variations in the conditional moments in (8), but as we shall see later on, the weights can facilitate such effects by allowing them to vary over time. A constant can trivially be included as one of the forecasts so that the combination scheme allows for an intercept term, a strategy recommended (under MSE loss) by Granger and Ramanathan (1984) and – for a more general class of loss functions – by Elliott and Timmermann (2004). Assuming that a constant is included, the optimal (population) values of the constant and the combination weights,  $\omega_{0t+h,t}^*$  and  $\omega_{t+h,t}^*$ , simplify as follows:

$$\omega_{0t+h,t}^* = \mu_{y_{t+h,t}} - \omega_{t+h,t}^{*\prime} \boldsymbol{\mu}_{\hat{y}_{t+h,t}}, \quad (9)$$

$$\omega_{t+h,t}^* = \boldsymbol{\Sigma}_{\hat{y}_{t+h,t}}^{-1} \boldsymbol{\sigma}_{y\hat{y}_{t+h,t}}.$$

These weights depend on the full conditional covariance matrix of the forecasts,  $\boldsymbol{\Sigma}_{\hat{y}_{t+h,t}}$ . In general the weights have an intuitive interpretation and tend to be larger for more accurate forecasts that are less strongly correlated with other forecasts. Notice that the constant,  $\omega_{0t+h,t}^*$ , corrects for any biases in the weighted forecast  $\omega_{t+h,t}^{*\prime} \hat{y}_{t+h,t}$ .

In the following we explore some interesting special cases to demonstrate the determinants of gains from forecast combination.

### 2.3.1. Diversification gains

Under quadratic loss it is easy to illustrate the population gains from different forecast combination schemes. This is an important task since, as argued by Winkler (1989, p. 607) “The better we understand which sets of underlying assumptions are associated with which combining rules, the more effective we will be at matching combining rules to forecasting situations.” To this end we consider the simple combination of two forecasts that give rise to errors  $e_1 = y - \hat{y}_1$  and  $e_2 = y - \hat{y}_2$ . Without risk of confusion we have dropped the time and horizon subscripts. Assuming that the individual forecast errors are unbiased, we have  $e_1 \sim (0, \sigma_1^2)$ ,  $e_2 \sim (0, \sigma_2^2)$  where  $\sigma_1^2 = \text{var}(e_1)$ ,  $\sigma_2^2 = \text{var}(e_2)$ ,  $\sigma_{12} = \rho_{12}\sigma_1\sigma_2$  is the covariance between  $e_1$  and  $e_2$  and  $\rho_{12}$  is their correlation. Suppose that the combination weights are restricted to sum to one, with weights  $(\omega, 1 - \omega)$  on the first and second forecast, respectively. The forecast error from the combination  $e^c = y - \omega\hat{y}_1 - (1 - \omega)\hat{y}_2$  takes the form

$$e^c = \omega e_1 + (1 - \omega)e_2. \quad (10)$$

By construction this has zero mean and variance

$$\sigma_c^2(\omega) = \omega^2\sigma_1^2 + (1 - \omega)^2\sigma_2^2 + 2\omega(1 - \omega)\sigma_{12}. \quad (11)$$

Differentiating with respect to  $\omega$  and solving the first order condition, we have

$$\omega^* = \frac{\sigma_2^2 - \sigma_{12}}{\sigma_1^2 + \sigma_2^2 - 2\sigma_{12}}, \quad (12)$$

$$1 - \omega^* = \frac{\sigma_1^2 - \sigma_{12}}{\sigma_1^2 + \sigma_2^2 - 2\sigma_{12}}.$$

A greater weight is assigned to models producing more precise forecasts (lower forecast error variances). A negative weight on a forecast clearly does not mean that it has no value to a forecaster. In fact when  $\rho_{12} > \sigma_2/\sigma_1$  the combination weights are not convex and one weight will exceed unity, the other being negative, cf. [Bunn \(1985\)](#).

Inserting  $\omega^*$  into the objective function (11), we get the expected squared loss associated with the optimal weights:

$$\sigma_c^2(\omega^*) = \frac{\sigma_1^2 \sigma_2^2 (1 - \rho_{12}^2)}{\sigma_1^2 + \sigma_2^2 - 2\rho_{12}\sigma_1\sigma_2}. \quad (13)$$

It can easily be verified that  $\sigma_c^2(\omega^*) \leq \min(\sigma_1^2, \sigma_2^2)$ . In fact, the diversification gain will only be zero in the following special cases (i)  $\sigma_1$  or  $\sigma_2$  equal to zero; (ii)  $\sigma_1 = \sigma_2$  and  $\rho_{12} = 1$ ; or (iii)  $\rho_{12} = \sigma_1/\sigma_2$ .

It is interesting to compare the variance of the forecast error from the optimal combination (12) to the variance of the combination scheme that weights the forecasts inversely to their relative mean squared error (MSE) values and hence ignores any correlation between the forecast errors:

$$\omega_{\text{inv}} = \frac{\sigma_2^2}{\sigma_1^2 + \sigma_2^2}, \quad 1 - \omega_{\text{inv}} = \frac{\sigma_1^2}{\sigma_1^2 + \sigma_2^2}. \quad (14)$$

These weights result in a forecast error variance

$$\sigma_{\text{inv}}^2 = \frac{\sigma_1^2 \sigma_2^2 (\sigma_1^2 + \sigma_2^2 + 2\rho_{12}\sigma_1\sigma_2)}{(\sigma_1^2 + \sigma_2^2)^2}. \quad (15)$$

After some algebra we can derive the ratio of the forecast error variance under this scheme relative to its value under the optimal weights,  $\sigma_c^2(\omega^*)$  in (13):

$$\frac{\sigma_{\text{inv}}^2}{\sigma_c^2(\omega^*)} = \left( \frac{1}{1 - \rho_{12}^2} \right) \left( 1 - \left( \frac{2\sigma_{12}}{\sigma_1^2 + \sigma_2^2} \right)^2 \right). \quad (16)$$

If  $\sigma_1 \neq \sigma_2$ , this exceeds unity unless  $\rho_{12} = 0$ . When  $\sigma_1 = \sigma_2$ , this ratio is always unity irrespective of the value of  $\rho_{12}$  and in this case  $\omega_{\text{inv}} = \omega^* = 1/2$ . Equal weights are optimal when combining two forecasts provided that the two forecast error variances are identical, irrespective of the correlation between the two forecast errors.

Another interesting benchmark is the equal-weighted combination  $\hat{y}^{ew} = (1/2)(\hat{y}_1 + \hat{y}_2)$ . Under these weights the variance of the forecast error is

$$\sigma_{ew}^2 = \frac{1}{4}\sigma_1^2 + \frac{1}{4}\sigma_2^2 + \frac{1}{2}\sigma_1\sigma_2\rho_{12} \tag{17}$$

so the ratio  $\sigma_{ew}^2/\sigma_c^2(\omega^*)$  becomes:

$$\frac{\sigma_{ew}^2}{\sigma_c^2(\omega^*)} = \frac{(\sigma_1^2 + \sigma_2^2)^2 - 4\sigma_{12}^2}{4\sigma_1^2\sigma_2^2(1 - \rho_{12}^2)}, \tag{18}$$

which in general exceeds unity unless  $\sigma_1 = \sigma_2$ .

Finally, as a measure of the diversification gain obtained from combining the two forecasts it is natural to compare  $\sigma_c^2(\omega^*)$  to  $\min(\sigma_1^2, \sigma_2^2)$ . Suppose that  $\sigma_1 > \sigma_2$  and define  $\kappa = \sigma_2/\sigma_1$  so that  $\kappa < 1$ . We then have

$$\frac{\sigma_c^2(\omega^*)}{\sigma_2^2} = \frac{1 - \rho_{12}^2}{1 + \kappa^2 - 2\rho_{12}\kappa}. \tag{19}$$

Figure 1 shows this expression graphically as a function of  $\rho_{12}$  and  $\kappa$ . The diversification gain is a complicated function of the correlation between the two forecast errors,  $\rho_{12}$ ,




Figure 1.

and the variance ratio of the forecast errors,  $\kappa$ . In fact, the derivative of the efficiency gain with respect to either  $\kappa$  or  $\rho_{12}$  changes sign even for reasonable parameter values. Differentiating (19) with respect to  $\rho_{12}$ , we have

$$\frac{\partial(\sigma_c^2(\omega^*)/\sigma_2^2)}{\partial\rho_{12}} \propto \kappa\rho_{12}^2 - (1 + \kappa^2)\rho_{12} + \kappa.$$

This is a second order polynomial in  $\rho_{12}$  with roots (assuming  $\kappa < 1$ )

$$\frac{1 + \kappa^2 \pm (1 - \kappa^2)}{2\kappa} = (\kappa; 1/\kappa).$$

Only when  $\kappa = 1$  (so  $\sigma_1^2 = \sigma_2^2$ ) does it follow that the efficiency gain will be an increasing function of  $\rho_{12}$  – otherwise it will change sign, being positive on the interval  $[-1; \kappa]$  and negative on  $[\kappa; 1]$  as can be seen from Figure 1. The figure shows that diversification through combination is more effective (in the sense that it results in the largest reduction in the forecast error variance for a given change in  $\rho_{12}$ ) when  $\kappa = 1$ .

### 2.3.2. Effect of bias in individual forecasts

Problems can arise for forecast combinations when one or more of the individual forecasts is biased, the combination weights are constrained to sum to unity and an intercept is omitted from the combination scheme. Min and Zellner (1993) illustrate how bias in one or more of the forecasts along with a constraint that the weights add up to unity can lead to suboptimality of combinations. Let  $y - \hat{y}_1 = e_1 \sim (0, \sigma^2)$  and  $y - \hat{y}_2 = e_2 \sim (\mu_2, \sigma^2)$ ,  $\text{cov}(e_1, e_2) = \sigma_{12} = \rho_{12}\sigma^2$ , so  $\hat{y}_1$  is unbiased while  $\hat{y}_2$  has a bias equal of  $\mu_2$ . Then the MSE of  $\hat{y}_1$  is  $\sigma^2$ , while the MSE of  $\hat{y}_2$  is  $\sigma^2 + \mu_2^2$ . The MSE of the combined forecast  $\hat{y}_c = \omega\hat{y}_1 + (1 - \omega)\hat{y}_2$  relative to that of the best forecast ( $\hat{y}_1$ ) is

$$\text{MSE}(\hat{y}_c) - \text{MSE}(\hat{y}_1) = (1 - \omega)\sigma^2 \left( (1 - \omega) \left( \frac{\mu_2}{\sigma} \right)^2 - 2\omega(1 - \rho_{12}) \right),$$

so  $\text{MSE}(\hat{y}_c) > \text{MSE}(\hat{y}_1)$  if

$$\left( \frac{\mu_2}{\sigma} \right)^2 > \frac{2\omega(1 - \rho_{12})}{1 - \omega}.$$

This condition always holds if  $\rho_{12} = 1$ . Furthermore, the larger the bias, the more likely it is that the combination will not dominate the first forecast. Of course the problem here is that the combination is based on variances and not the mean squared forecast errors which would account for the bias.

### 2.4. Optimality of equal weights – general case

Equally weighted combinations occupy a special place in the forecast combination literature. They are frequently either imposed on the combination scheme or used as a

point towards which the unconstrained combination weights are shrunk. Given their special role, it is worth establishing more general conditions under which they are optimal in a population sense. This sets a benchmark that proves helpful in understanding their good finite-sample performance in simulations and in empirical studies with actual data.

Let  $\Sigma_e = E[\mathbf{e}\mathbf{e}']$  be the covariance matrix of the individual forecast errors where  $\mathbf{e} = \boldsymbol{\iota}_y - \hat{\mathbf{y}}$  and  $\boldsymbol{\iota}$  is an  $N \times 1$  column vector of ones. Again we drop time and horizon subscripts without any risk of confusion. From (7) and assuming that the individual forecasts are unbiased, so  $\boldsymbol{\iota}'\hat{\boldsymbol{\mu}}_y = \boldsymbol{\mu}'_y \boldsymbol{\iota}'$ , the vector of forecast errors has second moment

$$\begin{aligned}\Sigma_e &= E[y^2 \boldsymbol{u}' + \hat{\mathbf{y}}\hat{\mathbf{y}}' - 2y\boldsymbol{\iota}\hat{\mathbf{y}}'] \\ &= (\sigma_y^2 + \mu_y^2)\boldsymbol{u}' + \boldsymbol{\mu}_y\boldsymbol{\mu}'_y + \Sigma_{\hat{\mathbf{y}}\hat{\mathbf{y}}} - 2\boldsymbol{\iota}\sigma'_{y\hat{\mathbf{y}}} - 2\mu_y\boldsymbol{\iota}\boldsymbol{\mu}'_y.\end{aligned}\quad (20)$$

Consider minimizing the expected forecast error variance subject to the constraint that the weights add up to one:

$$\begin{aligned}\min \quad & \boldsymbol{\omega}'\Sigma_e\boldsymbol{\omega} \\ \text{s.t.} \quad & \boldsymbol{\omega}'\boldsymbol{\iota} = 1.\end{aligned}\quad (21)$$

The constraint ensures unbiasedness of the combined forecast provided that  $\boldsymbol{\mu}'_y = \mu_y\boldsymbol{\iota}'$  so that

$$\mu_y^2\boldsymbol{u}' + \boldsymbol{\mu}_y\boldsymbol{\mu}'_y - 2\mu_y\boldsymbol{\iota}\boldsymbol{\mu}'_y = 0.$$

The Lagrangian associated with (21) is

$$\mathcal{L} = \boldsymbol{\omega}'\Sigma_e\boldsymbol{\omega} - \lambda(\boldsymbol{\omega}'\boldsymbol{\iota} - 1)$$

which yields the first order condition

$$\Sigma_e\boldsymbol{\omega} = \frac{\lambda}{2}\boldsymbol{\iota}.\quad (22)$$

Assuming that  $\Sigma_e$  is invertible, after pre-multiplying by  $\Sigma_e^{-1}\boldsymbol{\iota}'$  and recalling that  $\boldsymbol{\iota}'\boldsymbol{\omega} = 1$  we get  $\lambda/2 = (\boldsymbol{\iota}'\Sigma_e^{-1}\boldsymbol{\iota})^{-1}$ . Inserting this in (22) we have the frequently cited formula for the optimal weights:

$$\boldsymbol{\omega}^* = (\boldsymbol{\iota}'\Sigma_e^{-1}\boldsymbol{\iota})^{-1}\Sigma_e^{-1}\boldsymbol{\iota}.\quad (23)$$

Now suppose that the forecast errors have the same variance,  $\sigma^2$ , and correlation,  $\rho$ . Then we have

$$\begin{aligned}\Sigma_e^{-1} &= \frac{1}{\sigma^2(1-\rho)}\left(\mathbf{I} - \frac{\rho}{1+(N-1)\rho}\boldsymbol{u}'\right) \\ &= \frac{1}{\sigma^2(1-\rho)(1+(N-1)\rho)}((1+(N-1)\rho)\mathbf{I} - \rho\boldsymbol{u}'),\end{aligned}$$

where  $\mathbf{I}$  is the  $N \times N$  identity matrix. Inserting this in (23) we have

$$\begin{aligned}\boldsymbol{\Sigma}_e^{-1}\boldsymbol{t} &= \frac{\boldsymbol{t}}{\sigma^2(1+(N-1)\rho)}, \\ (\boldsymbol{t}'\boldsymbol{\Sigma}_e^{-1}\boldsymbol{t})^{-1} &= \frac{\sigma^2(1+(N-1)\rho)}{N},\end{aligned}$$

so

$$\boldsymbol{\omega}^* = \left(\frac{1}{N}\right)\boldsymbol{t}. \quad (24)$$

Hence equal-weights are optimal in situations with an arbitrary number of forecasts when the individual forecast errors have the same variance and identical pair-wise correlations. Notice that the property that the weights add up to unity only follows as a result of imposing the constraint  $\boldsymbol{t}'\boldsymbol{\omega} = 1$  and need not otherwise hold more generally.

### 2.5. Optimal combinations under asymmetric loss

Recent work has seen considerable interest in analyzing the effect of asymmetric loss on optimal predictions, cf., inter alia, Christoffersen and Diebold (1997), Granger and Pesaran (2000) and Patton and Timmermann (2004). These papers show that the standard properties of an optimal forecast under MSE loss case to hold under asymmetric loss. These properties include lack of bias, absence of serial correlation in the forecast error at the single-period forecast horizon and increasing forecast error variance as the horizon grows. It is therefore not surprising that asymmetric loss also affects combination weights. To illustrate the significance of the shape of the loss function for the optimal combination weights, consider linex loss. The linex loss function is convenient to use since it allows us to characterize the optimal forecast analytically. It takes the form, cf. Zellner (1986),

$$L(e_{t+h,t}) = \exp(ae_{t+h,t}) - ae_{t+h,t} + 1, \quad (25)$$

where  $a$  is a scalar that controls the aversion towards either positive ( $a > 0$ ) or negative ( $a < 0$ ) forecast errors and  $e_{t+h,t} = (y_{t+h} - \omega_{0t+h,t} - \boldsymbol{\omega}'_{t+h,t}\hat{\boldsymbol{y}}_{t+h,t})$ . First, suppose that the target variable and forecast are joint Gaussian with moments given in (7). Using the well-known result that if  $X \sim N(\mu, \sigma^2)$ , then  $E[e^x] = \exp(\mu + \sigma^2/2)$ , the optimal combination weights  $(\omega_{0t+h,t}^*, \boldsymbol{\omega}_{t+h,t}^*)$  which minimize the expected loss  $E[L(e_{t+h,t})|\mathcal{F}_t]$ , solve

$$\begin{aligned}\min_{\omega_{0t+h,t}, \boldsymbol{\omega}_{t+h,t}} \exp &\left( a(\mu_{y_{t+h,t}} - \omega_{0t+h,t} - \boldsymbol{\omega}'_{t+h,t}\boldsymbol{\mu}_{\hat{\boldsymbol{y}}_{t+h,t}}) \right. \\ &+ \frac{a^2}{2}(\sigma_{y_{t+h,t}}^2 + \boldsymbol{\omega}'_{t+h,t}\boldsymbol{\Sigma}_{\hat{\boldsymbol{y}}_{t+h,t}}\boldsymbol{\omega}_{t+h,t} - 2\boldsymbol{\omega}'_{t+h,t}\boldsymbol{\sigma}_{y_{t+h,t}}) \\ &\left. - a(\mu_{y_{t+h,t}} - \omega_{0t+h,t} - \boldsymbol{\omega}'_{t+h,t}\boldsymbol{\mu}_{\hat{\boldsymbol{y}}_{t+h,t}}) \right).\end{aligned}$$



Taking derivatives, we get the first order conditions

$$\begin{aligned} \exp\left(a\left(\mu_{y_{t+h,t}} - \omega_{0_{t+h,t}} - \boldsymbol{\omega}'_{t+h,t} \boldsymbol{\mu}_{\hat{y}_{t+h,t}}\right) \right. \\ \left. + \frac{a^2}{2}\left(\sigma_{y_{t+h,t}}^2 + \boldsymbol{\omega}'_{t+h,t} \boldsymbol{\Sigma}_{\hat{y}_{t+h,t}} \boldsymbol{\omega}_{t+h,t} - 2\boldsymbol{\omega}'_{t+h,t} \boldsymbol{\sigma}_{y_{\hat{y}_{t+h,t}}}\right)\right) = 1, \quad (26) \\ \left(-a\boldsymbol{\mu}_{\hat{y}_{t+h,t}} + \frac{a^2}{2}\left(2\boldsymbol{\Sigma}_{\hat{y}_{t+h,t}} \boldsymbol{\omega}_{t+h,t} - 2\boldsymbol{\sigma}_{y_{\hat{y}_{t+h,t}}}\right)\right) + a\boldsymbol{\mu}_{\hat{y}_{t+h,t}} = 0. \end{aligned}$$

It follows that  $\boldsymbol{\omega}_{t+h,t}^* = \boldsymbol{\Sigma}_{\hat{y}_{t+h,t}}^{-1} \boldsymbol{\sigma}_{y_{\hat{y}_{t+h,t}}}$  which when inserted in the first equation gives the optimal solution

$$\begin{aligned} \omega_{0_{t+h,t}}^* &= \mu_{y_{t+h,t}} - \boldsymbol{\omega}_{t+h,t}^{*'} \boldsymbol{\mu}_{\hat{y}_{t+h,t}} + \frac{a}{2}\left(\sigma_{y_{t+h,t}}^2 - \boldsymbol{\omega}_{t+h,t}^{*'} \boldsymbol{\sigma}_{y_{\hat{y}_{t+h,t}}}\right), \quad (27) \\ \boldsymbol{\omega}_{t+h,t}^* &= \boldsymbol{\Sigma}_{\hat{y}_{t+h,t}}^{-1} \boldsymbol{\sigma}_{y_{\hat{y}_{t+h,t}}}. \end{aligned}$$

Notice that the optimal combination weights,  $\boldsymbol{\omega}_{t+h,t}^*$ , are unchanged from the case with MSE loss, (9), while the intercept accounts for the shape of the loss function and depends on the parameter  $a$ . In fact, the optimal combination will have a bias,  $\frac{a}{2}(\sigma_{y_{t+h,t}}^2 - \boldsymbol{\omega}_{t+h,t}^{*'} \boldsymbol{\sigma}_{y_{\hat{y}_{t+h,t}}})$ , that reflects the dispersion of the forecast error evaluated at the optimal combination weights.

Next, suppose that we allow for a non-Gaussian forecast error distribution by assuming that the joint distribution of  $(y_{t+h}, \hat{y}_{t+h,t})'$  is a mixture of two Gaussian distributions driven by a state variable,  $S_{t+h}$ , which can take two values, i.e.  $s_{t+h} = 1$  or  $s_{t+h} = 2$  so that

$$\begin{pmatrix} y_{t+h} \\ \hat{y}_{t+h,t} \end{pmatrix} \sim N\left(\begin{pmatrix} \mu_{y_{s_{t+h}}} \\ \boldsymbol{\mu}_{\hat{y}_{s_{t+h}}} \end{pmatrix}, \begin{pmatrix} \sigma_{y_{s_{t+h}}}^2 & \boldsymbol{\sigma}'_{y_{\hat{y}_{s_{t+h}}}} \\ \boldsymbol{\sigma}_{y_{\hat{y}_{s_{t+h}}}} & \boldsymbol{\Sigma}_{\hat{y}_{s_{t+h}}} \end{pmatrix}\right). \quad (28)$$

Furthermore, suppose that  $P(S_{t+h} = 1) = p$ , while  $P(S_{t+h} = 2) = 1 - p$ . The two regimes could correspond to recession and expansion states for the economy [Hamilton (1989)] or bull and bear states for financial markets, cf. Guidolin and Timmermann (2005).

Under this model,

$$\begin{aligned} e_{t+h,t} &= y_{t+h} - \omega_{0_{t+h,t}} - \boldsymbol{\omega}'_{t+h,t} \hat{y}_{t+h,t} \\ &\sim N\left(\mu_{y_{s_{t+h}}} - \omega_{0_{t+h,t}} - \boldsymbol{\omega}'_{t+h,t} \boldsymbol{\mu}_{\hat{y}_{s_{t+h}}}, \sigma_{y_{s_{t+h}}}^2 + \boldsymbol{\omega}'_{t+h,t} \boldsymbol{\Sigma}_{\hat{y}_{s_{t+h}}} \boldsymbol{\omega}_{t+h,t} \right. \\ &\quad \left. - 2\boldsymbol{\omega}'_{t+h,t} \boldsymbol{\sigma}_{y_{\hat{y}_{s_{t+h}}}}\right). \end{aligned}$$

Dropping time and horizon subscripts, the expected loss under this distribution,  $E[L(e_{t+h,t})|\hat{y}_{t+h,t}]$ , is proportional to

$$\begin{aligned}
& p \left\{ \exp \left( a(\mu_{y1} - \omega_0 - \boldsymbol{\omega}' \boldsymbol{\mu}_{\hat{y}1}) + \frac{a^2}{2} (\sigma_{y1}^2 + \boldsymbol{\omega}' \boldsymbol{\Sigma}_{\hat{y}\hat{y}1} \boldsymbol{\omega} - 2\boldsymbol{\omega}' \boldsymbol{\sigma}_{y\hat{y}1}) \right) \right. \\
& \quad \left. - a(\mu_{y1} - \omega_0 - \boldsymbol{\omega}' \boldsymbol{\mu}_{\hat{y}1}) \right\} \\
& + (1-p) \left\{ \exp \left( a(\mu_{y2} - \omega_0 - \boldsymbol{\omega}' \boldsymbol{\mu}_{\hat{y}2}) + \frac{a^2}{2} (\sigma_{y2}^2 + \boldsymbol{\omega}' \boldsymbol{\Sigma}_{\hat{y}\hat{y}2} \boldsymbol{\omega} - 2\boldsymbol{\omega}' \boldsymbol{\sigma}_{y\hat{y}2}) \right) \right. \\
& \quad \left. - a(\mu_{y2} - \omega_0 - \boldsymbol{\omega}' \boldsymbol{\mu}_{\hat{y}2}) \right\}.
\end{aligned}$$

Taking derivatives, we get the following first order conditions for  $\omega_0$  and  $\boldsymbol{\omega}$ :

$$\begin{aligned}
& p(\exp(\xi_1) - 1) + (1-p)(\exp(\xi_2) - 1) = 0, \\
& p(\exp(\xi_1)(-\boldsymbol{\mu}_{\hat{y}1} + a(\boldsymbol{\Sigma}_{\hat{y}\hat{y}1} \boldsymbol{\omega} - \boldsymbol{\sigma}_{y\hat{y}1})) + \boldsymbol{\mu}_{\hat{y}1}) \\
& \quad + (1-p)(\exp(\xi_2)(-\boldsymbol{\mu}_{\hat{y}2} + a(\boldsymbol{\Sigma}_{\hat{y}\hat{y}2} \boldsymbol{\omega} - \boldsymbol{\sigma}_{y\hat{y}2})) + \boldsymbol{\mu}_{\hat{y}2}) = 0,
\end{aligned}$$

where

$$\xi_{s_{t+1}} = a(\mu_{y_{s_{t+1}}} - \omega_0 - \boldsymbol{\omega}' \boldsymbol{\mu}_{\hat{y}_{s_{t+1}}}) + \frac{a^2}{2} (\sigma_{y_{s_{t+1}}}^2 + \boldsymbol{\omega}' \boldsymbol{\Sigma}_{\hat{y}\hat{y}_{s_{t+1}}} \boldsymbol{\omega} - 2\boldsymbol{\omega}' \boldsymbol{\sigma}_{y\hat{y}_{s_{t+1}}}).$$

In general this gives a set of  $N+1$  highly nonlinear equations in  $\omega_0$  and  $\boldsymbol{\omega}$ . The exception is when  $\boldsymbol{\mu}_{\hat{y}1} = \boldsymbol{\mu}_{\hat{y}2}$ , in which case (using the first order condition for  $\omega_0$ ) the first order condition for  $\boldsymbol{\omega}$  simplifies to

$$p \exp(\xi_1)(\boldsymbol{\Sigma}_{\hat{y}\hat{y}1} \boldsymbol{\omega} - \boldsymbol{\sigma}_{y\hat{y}1}) + (1-p) \exp(\xi_2)(\boldsymbol{\Sigma}_{\hat{y}\hat{y}2} \boldsymbol{\omega} - \boldsymbol{\sigma}_{y\hat{y}2}) = \mathbf{0}.$$

When  $\boldsymbol{\Sigma}_{\hat{y}\hat{y}2} = \varphi \boldsymbol{\Sigma}_{\hat{y}\hat{y}1}$  and  $\boldsymbol{\sigma}_{y\hat{y}2} = \varphi \boldsymbol{\sigma}_{y\hat{y}1}$ , for any  $\varphi > 0$ , the solution to this equation again corresponds to the optimal weights for the MSE loss function, (9):

$$\boldsymbol{\omega}^* = \boldsymbol{\Sigma}_{\hat{y}\hat{y}1}^{-1} \boldsymbol{\sigma}_{y\hat{y}1}. \tag{29}$$

This restriction represents a very special case and ensures that the joint distribution of  $(y_{t+h}, \hat{y}_{t+h,t})$  is elliptically symmetric – a class of distributions that encompasses the multivariate Gaussian. This is a special case of the more general result by Elliott and Timmermann (2004): If the joint distribution of  $(y_{t+h}, \hat{y}'_{t+h,t})'$  is elliptically symmetric and the expected loss can be written as a function of the mean and variance of the forecast error,  $\mu_e$  and  $\sigma_e^2$ , i.e.,  $E[L(e_t)] = g(\mu_e, \sigma_e^2)$ , then the optimal forecast combination weights,  $\boldsymbol{\omega}^*$ , take the form (29) and hence do not depend on the shape of the loss function (other than for certain technical conditions). Conversely, the constant ( $\omega_0$ ) reflects this shape. Thus, under fairly general conditions on the loss functions, a forecast enters into the optimal combination with a non-zero weight if and only if its optimal weight under MSE loss is non-zero. Conversely, if elliptical symmetry fails to hold, then it is quite possible that a forecast may have a non-zero weight under loss functions

other than MSE loss but not under MSE loss and vice versa. The latter case is likely to be most relevant empirically since studies using regime switching models often find that, although the mean parameters may be constrained to be identical across regimes, the variance-covariance parameters tend to be very different across regimes, cf., e.g., **Guidolin and Timmermann (2005)**.

This example can be used to demonstrate that a forecast that adds value (in the sense that it is uncorrelated with the outcome variable) only a small part of the time when other forecasts break down, will be included in the optimal combination. We set all mean parameters equal to one,  $\mu_{y1} = \mu_{y2} = 1$ ,  $\boldsymbol{\mu}_{\hat{y}1} = \boldsymbol{\mu}_{\hat{y}2} = \boldsymbol{1}$ , so bias can be ignored, while the variance-covariance parameters are chosen as follows:

$$\begin{aligned} \sigma_{y1} &= 3; \quad \sigma_{y2} = 1, \\ \boldsymbol{\Sigma}_{\hat{y}\hat{y}1} &= 0.8 \times \sigma_{y1}^2 \times \mathbf{I}; \quad \boldsymbol{\Sigma}_{\hat{y}\hat{y}2} = 0.5 \times \sigma_{y2}^2 \times \mathbf{I}, \\ \sigma_{y\hat{y}1} &= \sigma_{y1} \times \sqrt{\text{diag}(\boldsymbol{\Sigma}_{\hat{y}\hat{y}1})} \odot \begin{pmatrix} 0.9 \\ 0.2 \end{pmatrix}, \\ \sigma_{y\hat{y}2} &= \sigma_{y2} \times \sqrt{\text{diag}(\boldsymbol{\Sigma}_{\hat{y}\hat{y}2})} \odot \begin{pmatrix} 0.0 \\ 0.8 \end{pmatrix}, \end{aligned}$$

where  $\odot$  is the Hadamard or element by element multiplication operator.

In **Table 1** we show the optimal weight on the two forecasts as a function of  $p$  for two different values of  $a$ , namely  $a = 1$ , corresponding to strongly asymmetric loss, and  $a = 0.1$ , representing less asymmetric loss. When  $p = 0.05$  and  $a = 1$ , so there is only a five percent chance that the process is in state 1, the optimal weight on model 1 is 35%. This is lowered to only 8% when the asymmetry parameter is reduced to  $a = 0.1$ . Hence the low probability event has a greater effect on the optimal combination weights the higher the degree of asymmetry in the loss function and the higher the variability of such events.

This example can also be used to demonstrate why forecast combinations may work when the underlying predictors are generated under different loss functions. Suppose that two forecasters have linex loss with parameters  $a_1 > 0$  and  $a_2 < 0$  and suppose

Table 1  
Optimal combination weights under asymmetric loss.

$a = 1$			$a = 0.1$		
$p$	$\omega_1^*$	$\omega_2^*$	$p$	$\omega_1^*$	$\omega_2^*$
0.05	0.346	0.324	0.05	0.081	0.365
0.10	0.416	0.314	0.10	0.156	0.353
0.25	0.525	0.297	0.25	0.354	0.323
0.50	0.636	0.280	0.50	0.620	0.283
0.75	0.744	0.264	0.75	0.831	0.250
0.90	0.842	0.249	0.90	0.940	0.234

that both have access to the same information set and use the same model to forecast the mean and variance of  $Y$ ,  $\hat{\mu}_{y_{t+h,t}}$ ,  $\hat{\sigma}_{y_{t+h,t}}^2$ . Their forecasts are then computed as [assuming normality, cf. Christoffersen and Diebold (1997)]

$$\hat{y}_{t+h,t,1} = \hat{\mu}_{y_{t+h,t}} + \frac{a_1}{2} \hat{\sigma}_{y_{t+h,t}}^2,$$

$$\hat{y}_{t+h,t,2} = \hat{\mu}_{y_{t+h,t}} + \frac{a_2}{2} \hat{\sigma}_{y_{t+h,t}}^2.$$

Each forecast includes an optimal bias whose magnitude is time-varying. For a forecast user with symmetric loss, neither of these forecasts is particularly useful as each is biased. Furthermore, the bias cannot simply be taken out by including a constant in the forecast combination regression since the bias is time-varying. However, in this simple case, there exists an exact linear combination of the two forecasts that is unbiased:

$$\hat{y}_{t+1,t}^c = \omega \hat{y}_{t+h,t,1} + (1 - \omega) \hat{y}_{t+h,t,2}, \quad \omega = \frac{-a_2}{a_1 - a_2}.$$

Of course this is a special case, but it nevertheless does show how biases in individual forecasts can either be eliminated or reduced in a forecast combination.

## 2.6. Combining as a hedge against non-stationarities

Hendry and Clements (2002) argue that forecast combinations may work well empirically because they provide insurance against what they refer to as extraneous (deterministic) structural breaks. They consider a wide array of simulation designs for the break and find that combinations work well under a shift in the intercept of a single variable in the data generating process. In addition when two or more positively correlated predictor variables are subject to shifts in opposite directions, forecast combinations can be expected to lead to even larger reductions in the MSE. Their analysis considers the case where a break occurs after the estimation period and does not affect the parameter estimates of the individual forecasting models. They establish conditions on the size of the post-sample break ensuring that an equal-weighted combination out-performs the individual forecasts.<sup>6</sup>

In support of the interpretation that structural breaks or model instability may explain the good average performance of forecast combination methods, Stock and Watson (2004) report that the performance of combined forecasts tends to be far more stable than that of the individual constituent forecasts entering in the combinations. Interestingly, however, many of the combination methods that attempt to build in time-variations in the combination weights (either in the form of discounting of past performance or time-varying parameters) have generally not proved to be successful, although there have been exceptions.

<sup>6</sup> See also Winkler (1989) who argues (p. 606) that "... in many situations there is no such thing as a 'true' model for forecasting purposes. The world around us is continually changing, with new uncertainties replacing old ones."

It is easy to construct examples of specific forms of non-stationarities in the underlying data generating process for which simple combinations work better than the forecast from the best single model. Aiolfi and Timmermann (2006) study the following simple model for changes or shifts in the data generating process:

$$\begin{aligned} y_t &= S_t f_{1t} + (1 - S_t) f_{2t} + \varepsilon_{yt}, \\ \hat{y}_{1t} &= f_{1t} + \varepsilon_{1t}, \\ \hat{y}_{2t} &= f_{2t} + \varepsilon_{2t}. \end{aligned} \quad (30)$$

All variables are assumed to be Gaussian with factors  $f_{1t} \sim N(\mu_1, \sigma_{f_1}^2)$ ,  $f_{2t} \sim N(\mu_2, \sigma_{f_2}^2)$  and innovations  $\varepsilon_{yt} \sim N(0, \sigma_{\varepsilon_y}^2)$ ,  $\varepsilon_{1t} \sim N(0, \sigma_{\varepsilon_1}^2)$ ,  $\varepsilon_{2t} \sim N(0, \sigma_{\varepsilon_2}^2)$ . Innovations are mutually uncorrelated and uncorrelated with the factors, while  $\text{Cov}(f_{1t}, f_{2t}) = \sigma_{f_1 f_2}$ . In addition, the state transition probabilities are constant:  $P(S_t = 1) = p$ ,  $P(S_t = 0) = 1 - p$ . Let  $\beta_1$  be the population projection coefficient of  $y_t$  on  $\hat{y}_{1t}$  while  $\beta_2$  is the population projection coefficient of  $y_t$  on  $\hat{y}_{2t}$ , so that

$$\begin{aligned} \beta_1 &= \frac{p\sigma_{f_1}^2 + (1-p)\sigma_{f_1 f_2}}{\sigma_{f_1}^2 + \sigma_{\varepsilon_1}^2}, \\ \beta_2 &= \frac{(1-p)\sigma_{f_2}^2 + p\sigma_{f_1 f_2}}{\sigma_{f_2}^2 + \sigma_{\varepsilon_2}^2}. \end{aligned}$$

The first and second moments of the forecast errors  $e_{it} = y_t - \hat{y}_{it}$ , can then be characterized as follows:

- Conditional on  $S_t = 1$ :

$$\begin{aligned} \begin{pmatrix} e_{1t} \\ e_{2t} \end{pmatrix} &\sim N\left(\begin{pmatrix} (1 - \beta_1)\mu_1 \\ \mu_1 - \beta_2\mu_2 \end{pmatrix}, \right. \\ &\quad \left. \begin{pmatrix} (1 - \beta_1)^2\sigma_{f_1}^2 + \beta_1^2\sigma_{\varepsilon_1}^2 + \sigma_{\varepsilon_y}^2 & (1 - \beta_1)\sigma_{f_1}^2 + \sigma_{\varepsilon_y}^2 \\ (1 - \beta_1)\sigma_{f_1}^2 + \sigma_{\varepsilon_y}^2 & \sigma_{f_1}^2 + \beta_2^2\sigma_{f_2}^2 + \beta_2^2\sigma_{\varepsilon_2}^2 + \sigma_{\varepsilon_y}^2 \end{pmatrix}\right). \end{aligned}$$

- Conditional on  $S_t = 0$ :

$$\begin{aligned} \begin{pmatrix} e_{1t} \\ e_{2t} \end{pmatrix} &\sim N\left(\begin{pmatrix} \mu_2 - \beta_1\mu_1 \\ (1 - \beta_2)\mu_2 \end{pmatrix}, \right. \\ &\quad \left. \begin{pmatrix} \beta_1^2\sigma_{f_1}^2 + \sigma_{f_2}^2 + \beta_1^2\sigma_{\varepsilon_1}^2 + \sigma_{\varepsilon_y}^2 & (1 - \beta_2)\sigma_{f_2}^2 + \sigma_{\varepsilon_y}^2 \\ (1 - \beta_2)\sigma_{f_2}^2 + \sigma_{\varepsilon_y}^2 & (1 - \beta_2)^2\sigma_{f_2}^2 + \beta_2^2\sigma_{\varepsilon_2}^2 + \sigma_{\varepsilon_y}^2 \end{pmatrix}\right). \end{aligned}$$

Under the joint model for  $(y_t, \hat{y}_{1t}, \hat{y}_{2t})$  in (30), Aiolfi and Timmermann (2006) show that the population MSE of the equal-weighted combined forecast will be lower than the population MSE of the best model provided that the following condition holds:

$$\frac{1}{3} \left( \frac{p}{1-p} \right)^2 \frac{1 + \psi_2}{1 + \psi_1} < \frac{\sigma_{f_2}^2}{\sigma_{f_1}^2} < 3 \left( \frac{p}{1-p} \right)^2 \frac{1 + \psi_2}{1 + \psi_1}. \quad (31)$$

Here  $\psi_1 = \sigma_{\varepsilon_1}^2 / \sigma_{f_1}^2$ ,  $\psi_2 = \sigma_{\varepsilon_2}^2 / \sigma_{f_2}^2$  are the noise-to-signal ratios for forecasts one and two, respectively. Hence if  $p = 1 - p = 1/2$  and  $\psi_1 = \psi_2$ , the condition in (31) reduces to

$$\frac{1}{3} < \frac{\sigma_{f_2}^2}{\sigma_{f_1}^2} < 3,$$

suggesting that equal-weighted combinations will provide a hedge against ‘breaks’ for a wide range of values of the relative factor variance. How good an approximation this model provides for actual data can be debated, but regime shifts have been widely documented for first and second moments of, *inter alia*, output growth, stock and bond returns, interest rates and exchange rates.

Conversely, when combination weights have to be estimated, instability in the data generating process may cause under-performance relative to that of the best individual forecasting model. Hence we can construct examples where combination is the dominant strategy in the absence of breaks or other forms of non-stationarities, but becomes inferior in the presence of breaks. This is likely to happen if the conditional distribution of the target variable given a particular forecast is stationary, whereas the correlations between the forecasts changes. In this case the combination weights will change but the individual models’ performance remain the same.

### 3. Estimation

Forecast combinations, while appealing in theory, are at a disadvantage over a single forecast model because they introduce parameter estimation error in cases where the combination weights need to be estimated. This is an important point – so much so, that seemingly suboptimal combination schemes such as equal-weighting have widely been found to dominate combination methods that would be optimal in the absence of parameter estimation errors. Finite-sample errors in the estimates of the combination weights can lead to poor performance of combination schemes that dominate in large samples.<sup>7</sup>

#### 3.1. To combine or not to combine

The first question to answer in the presence of multiple forecasts of the same variable is whether or not to combine the forecasts or rather simply attempt to identify the single

<sup>7</sup> Yang (2004) demonstrates theoretically that linear forecast combinations can lead to far worse performance than those from the best single forecasting model due to large variability in estimates of the combination weights and proposes a range of recursive methods for updating the combination weights that ensure that combinations achieve a performance similar to that of the best individual forecasting method up to a constant penalty term and a proportionality factor.

best forecasting model. Here it is important to distinguish between the situation where the information sets underlying the individual forecasts is observed from that where they are unobserved to the forecast user. When the information sets are unobserved it is often justified to combine forecasts provided that the private (non-overlapping) parts of the information sets are sufficiently important. Whether this is satisfied can be difficult to assess, but diagnostics such as the correlation between forecasts or forecast errors can be considered.

When forecast users do have access to the full information set used to construct the individual forecasts, [Chong and Hendry \(1986\)](#) and [Diebold \(1989\)](#) argue that combinations may be less justified. Successful combination indicates misspecification of the individual models and so a better individual model should be sought. Finding a ‘best’ model may of course be rather difficult if the space of models included in the search is high dimensional and the time-series short. As [Clemen \(1989\)](#) nicely puts it: “Using a combination of forecasts amounts to an admission that the forecaster is unable to build a properly specified model. Trying ever more elaborate combining models seems to add insult to injury as the more complicated combinations do not generally perform that well.”

Simple tests of whether one forecast dominates another forecast are neither sufficient nor necessary for settling the question of whether or not to combine. This follows since we can construct examples where (in population) forecast  $\hat{y}_1$  dominates forecast  $\hat{y}_2$  (in the sense that it leads to lower expected loss), yet it remains optimal to combine the two forecasts.<sup>8</sup> Similarly, we can construct examples where forecast  $\hat{y}_1$  and  $\hat{y}_2$  generate identical expected loss, yet it is not optimal to combine them – most obviously if they are perfectly correlated, but also due to estimation errors in the combination weights.

What is called for more generally is a test of whether one forecast – or a set of forecasts – encompasses all information contained in another forecast (or sets of forecasts). In the context of MSE loss functions, forecast encompassing tests have been developed by [Chong and Hendry \(1986\)](#). Point forecasts are sufficient statistics under MSE loss and a test of pair-wise encompassing can be based on the regression

$$y_{t+h} = \beta_0 + \beta_1 \hat{y}_{t+h,t,1} + \beta_2 \hat{y}_{t+h,t,2} + e_{t+h,t}, \quad t = 1, 2, \dots, T - h. \quad (32)$$

Forecast 1 encompasses forecast 2 when the parameter restriction  $(\beta_0 \ \beta_1 \ \beta_2) = (0 \ 1 \ 0)$  holds, while conversely if forecast 2 encompasses forecast 1 we have  $(\beta_0 \ \beta_1 \ \beta_2) = (0 \ 0 \ 1)$ . All other outcomes mean that there is some information in both forecasts which can then be usefully exploited. Notice that this is an argument that only holds in population. It is still possible in small samples that ignoring one forecast can lead to better out-of-sample forecasts even though, asymptotically, the coefficient on the omitted forecast in (32) differs from zero.

<sup>8</sup> Most obviously, under MSE loss, when  $\sigma(y - \hat{y}_1) > \sigma(y - \hat{y}_2)$ , and  $cor(y - \hat{y}_1, y - \hat{y}_2) \neq \sigma(y - \hat{y}_2)/\sigma(y - \hat{y}_1)$ , it will generally be optimal to combine the two forecasts, cf. Section 2.

More generally, a test that the forecast of some model, e.g., model 1, encompasses all other models can be based on a test of  $\beta_2 = \dots = \beta_N = 0$  in the regression

$$y_{t+h} - \hat{y}_{t+h,t,1} = \beta_0 + \sum_{i=2}^N \beta_i \hat{y}_{t+h,t,i} + e_{t+h,t}.$$

Inference is complicated by whether forecasting models are nested or non-nested, cf. West (2006), Chapter 3 in this Handbook, and the references therein.

In situations where the data is not very informative and it is not possible to identify a single dominant model, it makes sense to combine forecasts. Makridakis and Winkler (1983) explain this well (p. 990): “When a single method is used, the risk of not choosing the best method can be very serious. The risk diminishes rapidly when more methods are considered and their forecasts are averaged. In other words, the choice of the best method or methods becomes less important when averaging.” They demonstrate this point by showing that the forecasting performance of a combination strategy improves as a function of the number of models involved in the combination, albeit at a decreasing rate.

Swanson and Zeng (2001) propose to use model selection criteria such as the SIC to choose which subset of forecasts to combine. This approach does not require formal hypothesis testing so that size distortions due to the use of sequential pre-tests, can be avoided. Of course, consistency of the selection approach must be established in the context of the particular sampling experiment appropriate for a given forecasting situation. In empirical work reported by these authors the combination chosen by SIC appears to provide the best overall performance and rarely gets dominated by other methods in out-of-sample forecasting experiments.

Once it has been established whether to combine or not, there are various ways in which the combination weights,  $\hat{\omega}_{t+h,t}$ , can be estimated. We will discuss some of these methods in what follows. A theme that is common across estimators is that estimation errors in forecast combinations are generally important especially in cases where the number of forecasts,  $N$ , is large relative to the length of the time-series,  $T$ .

### 3.2. Least squares estimators of the weights

It is common to assume a linear-in-weights model and estimate combination weights by ordinary least squares, regressing realizations of the target variable,  $y_\tau$  on the  $N$ -vector of forecasts,  $\hat{y}_\tau$  using data over the period  $\tau = h, \dots, t$ :

$$\hat{\omega}_{t+h,t} = \left( \sum_{\tau=1}^{t-h} \hat{y}_{\tau+h,\tau} \hat{y}'_{\tau+h,\tau} \right)^{-1} \sum_{\tau=1}^{t-h} \hat{y}_{\tau+h,\tau} y_{\tau+h}. \quad (33)$$

Different versions of this basic least squares projection have been proposed. Granger and Ramanathan (1984) consider three regressions



$$\begin{aligned}
\text{(i)} \quad & y_{t+h} = \omega_{0h} + \boldsymbol{\omega}'_h \hat{\mathbf{y}}_{t+h,t} + \varepsilon_{t+h}, \\
\text{(ii)} \quad & y_{t+h} = \boldsymbol{\omega}'_h \hat{\mathbf{y}}_{t+h,t} + \varepsilon_{t+h}, \\
\text{(iii)} \quad & y_{t+h} = \boldsymbol{\omega}'_h \hat{\mathbf{y}}_{t+h,t} + \varepsilon_{t+h}, \quad \text{s.t.} \quad \boldsymbol{\omega}'_h \mathbf{1} = 1.
\end{aligned} \tag{34}$$

The first and second of these regressions can be estimated by standard least squares, the only difference being that the second equation omits an intercept term. The third regression omits an intercept and can be estimated through constrained least squares. The first, and most general, regression does not require that the individual forecasts are unbiased since any bias can be adjusted through the intercept term,  $\omega_{0h}$ . In contrast, the third regression is motivated by an assumption of unbiasedness of the individual forecasts. Imposing that the weights sum to one then guarantees that the combined forecast is also unbiased. This specification may not be efficient, however, as the latter constraint can lead to efficiency losses as  $E[\hat{\mathbf{y}}_{t+h,t} \varepsilon_{t+h}] \neq \mathbf{0}$ . One could further impose convexity constraints  $0 \leq \omega_{h,i} \leq 1$ ,  $i = 1, \dots, N$ , to rule out that the combined forecast lies outside the range of the individual forecasts.

Another reason for imposing the constraint  $\boldsymbol{\omega}'_h \mathbf{1} = 1$  has been discussed by Diebold (1988). He proposes the following decomposition of the forecast error from the combination regression:

$$\begin{aligned}
e_{t+h,t}^c &= y_{t+h} - \omega_{0h} - \boldsymbol{\omega}'_h \hat{\mathbf{y}}_{t+h,t} \\
&= -\omega_{0h} + (1 - \boldsymbol{\omega}'_h \mathbf{1}) y_{t+h} + \boldsymbol{\omega}'_h (y_{t+h} \mathbf{1} - \hat{\mathbf{y}}_{t+h,t}) \\
&= -\omega_{0h} + (1 - \boldsymbol{\omega}'_h \mathbf{1}) y_{t+h} + \boldsymbol{\omega}'_h \mathbf{e}_{t+h,t},
\end{aligned} \tag{35}$$

where  $\mathbf{e}_{t+h,t}$  is the  $N \times 1$  vector of  $h$ -period forecast errors from the individual models. Oftentimes the target variable,  $y_{t+h}$ , is quite persistent whereas the forecast errors from the individual models are not serially correlated even when  $h = 1$ . It follows that unless it is imposed that  $1 - \boldsymbol{\omega}'_h \mathbf{1} = 0$ , then the forecast error from the combination regression typically will be serially correlated and hence be predictable itself.

### 3.3. Relative performance weights

Estimation errors in the combination weights tend to be particularly large due to difficulties in precisely estimating the covariance matrix,  $\boldsymbol{\Sigma}_e$ . One answer to this problem is to simply ignore correlations across forecast errors. Combination weights that reflect the performance of each individual model relative to the performance of the average model, but ignore correlations across forecasts have been proposed by Bates and Granger (1969) and Newbold and Granger (1974). Both papers argue that correlations can be poorly estimated and should be ignored in situations with many forecasts and short time-series. This effectively amounts to treating  $\boldsymbol{\Sigma}_e$  as a diagonal matrix, cf. Winkler and Makridakis (1983).

Stock and Watson (2001) propose a broader set of combination weights that also ignore correlations between forecast errors but base the combination weights on the models' relative MSE performance raised to various powers. Let  $\text{MSE}_{t+h,t,i} =$

$(1/v) \sum_{\tau=t-v}^t e_{\tau,\tau-h,i}^2$  be the  $i$ th forecasting model's MSE at time  $t$ , computed over a window of the previous  $v$  periods. Then

$$\hat{y}_{t+h,t}^c = \sum_{i=1}^N \hat{\omega}_{t+h,t,i} \hat{y}_{t+h,t,i}, \quad \hat{\omega}_{t+h,t,i} = \frac{(1/\text{MSE}_{t+h,t,i}^\kappa)}{\sum_{j=1}^N (1/\text{MSE}_{t+h,t,j}^\kappa)}. \quad (36)$$

Setting  $\kappa = 0$  assigns equal weights to all forecasts, while forecasts are weighted by the inverse of their MSE when  $\kappa = 1$ . The latter strategy has been found to work well in practice as it does not require estimating the off-diagonal parameters of the covariance matrix of the forecast errors. Such weights therefore disregard any correlations between forecast errors and so are only optimal in large samples provided that the forecast errors are truly uncorrelated.

### 3.4. Moment estimators

Outside the quadratic loss framework one can base estimation of the combination weights directly on the loss function, cf. Elliott and Timmermann (2004). Let the realized loss in period  $t + h$  be

$$L(e_{t+h,t}; \omega) = L(\omega | y_{t+h}, \hat{y}_{t+h,t}, \psi_L),$$

where  $\psi_L$  are the (given) parameters of the loss function. Then  $\tilde{\omega}_h = (\omega_{0h} \omega'_h)'$  can be obtained as an  $M$ -estimator based on the sample analog of  $E[L(e_{t+h,t})]$  using a sample of  $T - h$  observations  $\{y_\tau, \hat{y}_{\tau,\tau-h}\}_{\tau=h+1}^T$ :

$$\bar{L}(\omega) = (T - h)^{-1} \sum_{\tau=h+1}^T L(e_{\tau,\tau-h}(\tilde{\omega}_h); \theta_L).$$

Taking derivatives, one can use the generalized method of moments (GMM) to estimate  $\omega_{T+h,T}$  from the quadratic form

$$\min_{\tilde{\omega}_h} \left( \sum_{\tau=h+1}^T \mathbf{L}'(e_{\tau,\tau-h}(\tilde{\omega}_h); \psi_L) \right)' \mathbf{\Lambda}^{-1} \left( \sum_{\tau=h+1}^T \mathbf{L}'(e_{\tau,\tau-h}(\tilde{\omega}_h); \psi_L) \right), \quad (37)$$

where  $\mathbf{\Lambda}$  is a (positive definite) weighting matrix and  $\mathbf{L}'$  is a vector of derivatives of the moment conditions with respect to  $\tilde{\omega}_h$ . Consistency and asymptotic normality of the estimated weights is easily established under standard regularity conditions.

### 3.5. Nonparametric combination schemes

The estimators considered so far require stationarity at least for the moments involved in the estimation. To be empirically successful, they also require a reasonably large data sample (relative to the number of models,  $N$ ) as they otherwise tend not to be robust to outliers, cf. Gupta and Wilton (1987, p. 358): "... combination weights derived using

minimum variance or regression are not robust given short data samples, instability or nonstationarity. This leads to poor performance in the prediction sample.” In many applications the number of forecasts,  $N$ , is large relatively to the length of the time-series,  $T$ . In this case, it is not feasible to estimate the combination weights by OLS. Simple combination schemes such as an equal-weighted average of forecasts  $y_{t+h,t}^{\text{ew}} = t' \hat{\mathbf{y}}_{t+h,t} / N$  or weights based on the inverse MSE-values offer are an attractive option in this situation.

Simple, rank-based weighting schemes can also be constructed and have been used with some success in mean-variance analysis in finance, cf. [Wright and Satchell \(2003\)](#). These take the form  $\omega_{t+h,t} = f(\mathcal{R}_{t,t-h,1}, \dots, \mathcal{R}_{t,t-h,N})$ , where  $\mathcal{R}_{t,t-h,i}$  is the rank of the  $i$ th model based on its  $h$ -period performance up to time  $t$ . The most common scheme in this class is to simply use the median forecast as proposed by authors such as [Armstrong \(1989\)](#), [Hendry and Clements \(2002\)](#) and [Stock and Watson \(2001, 2004\)](#). Alternatively one can consider a triangular weighting scheme that lets the combination weights be inversely proportional to the models' rank, cf. [Aiolfi and Timmermann \(2006\)](#):

$$\hat{\omega}_{t+h,t,i} = \mathcal{R}_{t,t-h,i}^{-1} / \left( \sum_{i=1}^N \mathcal{R}_{t,t-h,i}^{-1} \right). \quad (38)$$

Again this combination ignores correlations across forecast errors. However, since ranks are likely to be less sensitive to outliers, this weighting scheme can be expected to be more robust than the weights in (33) or (36).

Another example in this class is spread combinations. These have been proposed by [Aiolfi and Timmermann \(2006\)](#) and consider weights of the form

$$\hat{\omega}_{t+h,t,i} = \begin{cases} \frac{1 + \bar{\omega}}{\alpha N} & \text{if } \mathcal{R}_{t,t-h,i} \leq \alpha N, \\ 0 & \text{if } \alpha N < \mathcal{R}_{t,t-h,i} < (1 - \alpha)N, \\ \frac{-\bar{\omega}}{\alpha N} & \text{if } \mathcal{R}_{t,t-h,i} \leq (1 - \alpha)N, \end{cases} \quad (39)$$

where  $\alpha$  is the proportion of top models that – based on performance up to time  $t$  – gets a weight of  $(1 + \bar{\omega})/\alpha N$ . Similarly, a proportion  $\alpha$  of models gets a weight of  $-\bar{\omega}/\alpha N$ . The larger the value of  $\alpha$ , the wider the set of top and bottom models that are used in the combination. Similarly, the larger is  $\bar{\omega}$ , the bigger the difference in weights on top and bottom models. The intuition for such spread combinations can be seen from (12) when  $N = 2$  so  $\alpha = 1/2$ . Solving for  $\rho_{12}$  we see that  $\omega^* = 1 + \bar{\omega}$  provided that

$$\rho_{12} = \frac{1}{2\bar{\omega} + 1} \left( \frac{\sigma_2}{\sigma_1} \bar{\omega} + \frac{\sigma_1}{\sigma_2} (1 + \bar{\omega}) \right).$$

Hence if  $\sigma_1 \approx \sigma_2$ , spread combinations are close to optimal provided that  $\rho_{12} \approx 1$ . The second forecast provides a hedge for the performance of the first forecast in this

situation. In general, spread portfolios are likely to work well when the forecasts are strongly collinear.

Gupta and Wilton (1987) propose an odds ratio combination approach based on a matrix of pair-wise odds ratios. Let  $\pi_{ij}$  be the probability that the  $i$ th forecasting model outperforms the  $j$ th model out-of-sample. The ratio  $o_{ij} = \pi_{ij}/\pi_{ji}$  is then the odds that model  $i$  will outperform model  $j$  and  $o_{ij} = 1/o_{ji}$ . Filling out the  $N \times N$  odds ratio matrix  $O$  with  $i, j$  element  $o_{ij}$  requires specifying  $N(N - 1)/2$  pairs of probabilities of outperformance,  $\pi_{ij}$ . An estimate of the combination weight  $\omega$  is obtained from the solution to the system of equations  $(\mathbf{O} - N\mathbf{I})\omega = \mathbf{0}$ . Since  $\mathbf{O}$  has unit rank with a trace equal to  $N$ ,  $\omega$  can be found as the normalized eigenvector associated with the largest (and only non-zero) eigenvalue of  $\mathbf{O}$ . This approach gives weights that are insensitive to small changes in the odds ratio and so does not require large amounts of data. Also, as it does not account for dependencies between the models it is likely to be less sensitive to changes in the covariance matrix than the regression approach. Conversely, it can be expected to perform worse if such correlations are important and can be estimated with sufficient precision.<sup>9</sup>

### 3.6. Pooling, clustering and trimming

Rather than combining the full set of forecasts, it is often advantageous to discard the models with the worst performance (trimming). Combining only the best models goes under the header ‘use sensible models’ in Armstrong (1989). This is particularly important when forecasting with nonlinear models whose predictions are often implausible and can lie outside the empirical range of the target variable. One can base whether or not to trim – and by how much to trim – on formal tests or on more loose decision rules.

To see why trimming can be important, suppose a fraction  $\alpha$  of the forecasting models contain valuable information about the target variable while a fraction  $1 - \alpha$  is pure noise. It is easy to see in this extreme case that the optimal forecast combination puts zero weight on the pure noise forecasts. However, once combination weights have to be estimated, forecasts that only add marginal information should be dropped from the combination since the cost of their inclusion – increased parameter estimation error – is not matched by similar benefits.

<sup>9</sup> Bunn (1975) proposes a combination scheme with weights reflecting the probability that a model produces the lowest loss, i.e.

$$p_{t+h,t,i} = \Pr(L(e_{t+h,t,i}) < L(e_{t+h,t,j})) \quad \text{for all } j \neq i,$$

$$\hat{y}_{t+h,t}^c = \sum_{i=1}^N p_{t+h,t,i} \hat{y}_{t+h,t,i}.$$

Bunn discusses how  $p_{t+h,t,i}$  can be updated based on a model’s track historical record using the proportion of times up to the current period where a model outperformed its competitors.

The ‘thick modeling’ approach – thus named because it seeks to exploit information in a cross-section (thick set) of models – proposed by Granger and Jeon (2004) is an example of a trimming scheme that removes poorly performing models in a step that precedes calculation of combination weights. Granger and Jeon argue that “an advantage of thick modeling is that one no longer needs to worry about difficult decisions between close alternatives or between deciding the outcome of a test that is not decisive”.

Grouping or clustering of forecasts can be motivated by the assumption of a common factor structure underlying the forecasting models. Consider the factor model

$$Y_{t+h} = \mu_y + \boldsymbol{\beta}'_y \mathbf{f}_{t+h} + \varepsilon_{yt+h},$$

$$\hat{y}_{t+h,t} = \mu_{\hat{y}} + \mathbf{B} \mathbf{f}_{t+h} + \boldsymbol{\varepsilon}_{t+h}, \quad (40)$$

where  $\mathbf{f}_{t+h}$  is an  $n_f \times 1$  vector of factor realizations satisfying  $E[\mathbf{f}_{t+h} \varepsilon_{yt+h}] = \mathbf{0}$ ,  $E[\mathbf{f}_{t+h} \boldsymbol{\varepsilon}'_{t+h}] = \mathbf{0}$  and  $E[\mathbf{f}_{t+h} \mathbf{f}'_{t+h}] = \boldsymbol{\Sigma}_f$ .  $\boldsymbol{\beta}_y$  is an  $n_f \times 1$  vector while  $\mathbf{B}$  is an  $N \times n_f$  matrix of factor loadings. For simplicity we assume that the factors have been orthogonalized. This will obviously hold if they are constructed as the principal components from a large data set and can otherwise be achieved through rotation. Furthermore, all innovations  $\varepsilon$  are serially uncorrelated with zero mean,  $E[\varepsilon^2_{yt+h}] = \sigma_{\varepsilon_y}^2$ ,  $E[\varepsilon_{yt+h} \boldsymbol{\varepsilon}_{t+h}] = \mathbf{0}$  and the noise in the individual forecasts is assumed to be idiosyncratic (model specific), i.e.,

$$E[\varepsilon_{it+h} \varepsilon_{jt+h}] = \begin{cases} \sigma_{\varepsilon_i}^2 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

We arrange these values on a diagonal matrix  $E[\boldsymbol{\varepsilon}_{t+h} \boldsymbol{\varepsilon}'_{t+h}] = \mathbf{D}_\varepsilon$ . This gives the following moments:

$$\begin{pmatrix} y_{t+h} \\ \hat{y}_{t+h,t} \end{pmatrix} \sim \begin{pmatrix} \mu_y \\ \mu_{\hat{y}} \end{pmatrix}, \begin{pmatrix} \boldsymbol{\beta}'_y \boldsymbol{\Sigma}_f \boldsymbol{\beta}_y + \sigma_{\varepsilon_y}^2 & \boldsymbol{\beta}'_y \boldsymbol{\Sigma}_f \mathbf{B}' \\ \mathbf{B} \boldsymbol{\Sigma}_f \boldsymbol{\beta}_y & \mathbf{B} \boldsymbol{\Sigma}_f \mathbf{B}' + \mathbf{D}_\varepsilon \end{pmatrix}.$$

Also suppose either that  $\mu_{\hat{y}} = \mathbf{0}$ ,  $\mu_y = 0$  or a constant is included in the combination scheme. Then the first order condition for the optimal weights is, from (8),

$$\boldsymbol{\omega}^* = (\mathbf{B} \boldsymbol{\Sigma}_f \mathbf{B}' + \mathbf{D}_\varepsilon)^{-1} \mathbf{B} \boldsymbol{\Sigma}_f \boldsymbol{\beta}_y. \quad (41)$$

Further suppose that the  $N$  forecasts of the  $n_f$  factors can be divided into appropriate groups according to their factor loading vectors  $\mathbf{b}_i$  such that  $\sum_{i=1}^{n_f} \dim(\mathbf{b}_i) = N$ :

$$\mathbf{B} = \begin{pmatrix} \mathbf{b}_1 & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{b}_2 & \mathbf{0} & \dots \\ \vdots & \mathbf{0} & \ddots & \mathbf{0} \\ \mathbf{0} & \dots & \mathbf{0} & \mathbf{b}_{n_f} \end{pmatrix}.$$

Then the first term on the right-hand side of (41) is given by

$$\mathbf{B}\Sigma_f\mathbf{B}' + \mathbf{D}_\varepsilon = \begin{pmatrix} \mathbf{b}_1\mathbf{b}'_1 & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{b}_2\mathbf{b}'_2 & \mathbf{0} & \dots \\ \vdots & \mathbf{0} & \ddots & \mathbf{0} \\ \mathbf{0} & \dots & \mathbf{0} & \mathbf{b}_{n_f}\mathbf{b}'_{n_f} \end{pmatrix} \mathbf{D}_{\sigma_F^2} + \mathbf{D}_\varepsilon, \quad (42)$$

where  $\mathbf{D}_{\sigma_F}$  is a diagonal matrix with  $\sigma_{f_1}^2$  in its first  $n_1$  diagonal places followed by  $\sigma_{f_2}^2$  in the next  $n_2$  diagonal places and so on and  $\mathbf{D}_\varepsilon$  is a diagonal matrix with  $\text{Var}(\varepsilon_{it})$  as the  $i$ th diagonal element. Thus the matrix in (42) and its inverse will be block diagonal. Provided that the forecasts tracking the individual factors can be grouped and have similar factor exposure ( $\mathbf{b}_i$ ) within each group, this suggests that little is lost by pooling forecasts within each cluster and ignoring correlations across clusters. In a subsequent step, sample counterparts of the optimal combination weights for the grouped forecasts can be obtained by least-squares estimation. In this way, far fewer combination weights ( $n_f$  rather than  $N$ ) have to be estimated. This can be expected to decrease forecast errors and thus improve forecasting performance.

Building on these ideas Aiolfi and Timmermann (2006) propose to sort forecasting models into clusters using a K-mean clustering algorithm based on their past MSE performance. As the previous argument suggests, one could alternatively base clustering on correlation patterns among the forecast errors.<sup>10</sup> Their method identifies  $K$  clusters. Let  $\hat{\mathbf{y}}_{t+h,t}^k$  be the  $p_k \times 1$  vector containing the subset of forecasts belonging to cluster  $k$ ,  $k = 1, 2, \dots, K$ . By ordering the clusters such that the first cluster contains models with the lowest historical MSE values, Aiolfi and Timmermann consider three separate strategies. The first simply computes the average forecast across models in the cluster of previous best models:

$$\hat{\mathbf{y}}_{t+h,t}^{\text{CPB}} = (\mathbf{1}'_{p_1}/p_1)\hat{\mathbf{y}}_{t+h,t}^1. \quad (43)$$

A second combination strategy identifies a small number of clusters, pools forecasts within each cluster and then estimates optimal weights on these pooled predictions by least squares:

$$\hat{\mathbf{y}}_{t+h,t}^{\text{CLS}} = \sum_{k=1}^K \hat{\omega}_{t+h,t,k} [(\mathbf{1}'_{p_k}/p_k)\hat{\mathbf{y}}_{t+h,t}^k], \quad (44)$$

where  $\hat{\omega}_{t+h,t,k}$  are least-squares estimates of the optimal combination weights for the  $K$  clusters. This strategy is likely to work well if the variation in forecasting performance within each cluster is small relative to the variation in forecasting performance across clusters.

<sup>10</sup> The two clustering methods will be similar if  $\sigma_{F_i}$  varies significantly across factors and the factor exposure vectors,  $\mathbf{b}_i$ , and error variances  $\sigma_{\varepsilon_i}^2$  are not too dissimilar across models. In this case forecast error variances will tend to cluster around the factors that the various forecasting models are most exposed to.

Finally, the third strategy pools forecasts within each cluster, estimates least squares combination weights and then shrinks these towards equal weights in order to reduce the effect of parameter estimation error

$$\hat{y}_{t+h,t}^{\text{CSW}} = \sum_{k=1}^K \hat{s}_{t+h,t,k} [(\mathbf{l}'_{p_k} / p_k) \hat{\mathbf{y}}_{t+h,t}^k],$$

where  $\hat{s}_{t+h,t,k}$  are the shrinkage weights for the  $K$  clusters computed as  $\hat{s}_{t+h,t,k} = \lambda \hat{\omega}_{t+h,t,k} + (1 - \lambda) \frac{1}{K}$ ,  $\lambda = \max\{0, 1 - \kappa(\frac{K}{t-h-K})\}$ . The higher is  $\kappa$ , the higher the shrinkage towards equal weights.

#### 4. Time-varying and nonlinear combination methods

So far our analysis has concentrated on forecast combination schemes that assumed constant and linear combination weights. While this follows naturally in the case with MSE loss and a time-invariant Gaussian distribution for the forecasts and realization, outside this framework it is natural to consider more general combination schemes. Two such families of special interest that generalize (6) are linear combinations with time-varying weights:

$$\hat{y}_{t+h,t}^c = \omega_{0t+h,t} + \boldsymbol{\omega}'_{t+h,t} \hat{\mathbf{y}}_{t+h,t}, \tag{45}$$

where  $\omega_{0t+h,t}$ ,  $\boldsymbol{\omega}'_{t+h,t}$  are adapted to  $\mathcal{F}_t$ , and nonlinear combinations with constant weights:

$$\hat{y}_{t+h,t}^c = C(\hat{\mathbf{y}}_{t+h,t}, \boldsymbol{\omega}), \tag{46}$$

where  $C(\cdot)$  is some function that is nonlinear in the parameters,  $\boldsymbol{\omega}$ , in the vector of forecasts,  $\hat{\mathbf{y}}_{t+h,t}$ , or in both. There is a close relationship between time-varying and nonlinear combinations. For example, nonlinearities in the true data generating process can lead to time-varying covariances for the forecast errors and hence time-varying weights in the combination of (misspecified) forecasts.

We next describe some of the approaches within these classes that have been proposed in the literature.

##### 4.1. Time-varying weights

When the joint distribution of  $(y_{t+h} \hat{\mathbf{y}}'_{t+h,t})'$  – or at least its first and second moments – vary over time, it can be beneficial to let the combination weights change over time. Indeed, [Bates and Granger \(1969\)](#) and [Newbold and Granger \(1974\)](#) suggested either assigning a disproportionately large weight to the model that has performed best most recently or using an adaptive updating scheme that puts more emphasis on recent performance in assigning the combination weights. Rather than explicitly modeling the structure of the time-variation in the combination weights, [Bates and Granger](#) proposed

five adaptive estimation schemes based on exponential discounting or the use of rolling estimation windows.

The first combination scheme uses a rolling window of the most recent  $v$  observations based on the forecasting models' relative performance<sup>11</sup>

$$\hat{\omega}_{t,t-h,i}^{\text{BG1}} = \frac{(\sum_{\tau=t-v+1}^t e_{\tau,\tau-h,i}^2)^{-1}}{\sum_{j=1}^N (\sum_{\tau=t-v+1}^t e_{\tau,\tau-h,j}^2)^{-1}}. \quad (47)$$

The shorter is  $v$ , the more weight is put on the models' recent track record and the larger the part of the historical data that is discarded. If  $v = t$ , an expanding window is used and this becomes a special case of (36). Correlations between forecast errors are ignored by this scheme.

The second rolling window scheme accounts for such correlations across forecast errors but, again, only uses the most recent  $v$  observations for estimation:

$$\hat{\omega}_{t,t-h}^{\text{BG2}} = \frac{\hat{\Sigma}_{et,t-h}^{-1}}{t' \hat{\Sigma}_{et,t-h}^{-1} t}, \quad (48)$$

$$\hat{\Sigma}_{et,t-h}[i, j] = v^{-1} \sum_{\tau=t-v+1}^t e_{\tau,\tau-h,i} e_{\tau,\tau-h,j}.$$

The third combination scheme uses adaptive updating captured by the parameter  $\alpha \in (0, 1)$ , which tends to smooth the time-series evolution in the combination weights:

$$\hat{\omega}_{t,t-h,i}^{\text{BG3}} = \alpha \hat{\omega}_{t-1,t-h-1,i} + (1 - \alpha) \frac{(\sum_{\tau=t-v+1}^t e_{\tau,\tau-h,i}^2)^{-1}}{\sum_{j=1}^N (\sum_{\tau=t-v+1}^t e_{\tau,\tau-h,j}^2)^{-1}}. \quad (49)$$

The closer to unity is  $\alpha$ , the smoother the weights will generally be.

The fourth and fifth combination methods are based on exponential discounting versions of the first two methods and take the form

$$\hat{\omega}_{t,t-h,i}^{\text{BG4}} = \frac{(\sum_{\tau=1}^t \lambda^\tau e_{\tau,\tau-h,i}^2)^{-1}}{\sum_{j=1}^N (\sum_{\tau=1}^t \lambda^\tau e_{\tau,\tau-h,j}^2)^{-1}}, \quad (50)$$

where  $\lambda \geq 1$  and higher values of  $\lambda$  correspond to putting more weight on recent data. This scheme does not put a zero weight on any of the past forecast errors whereas the rolling window methods entirely ignore observations more than  $v$  periods old. If  $\lambda = 1$ , there is no discounting of past performance and the formula becomes a special case of (36). However, it is common to use a discount factor such as  $\lambda = 1.05$  or  $\lambda = 1.10$ , although the chosen value will depend on factors such as data frequency, evidence of instability, forecast horizon, etc.

<sup>11</sup> While we write the equations for the weights for general  $h$ , adjustments can be made when  $h \geq 2$  which induces serial correlation in the forecast errors.



Finally, the fifth scheme estimates the variance and covariance of the forecast errors using exponential discounting:

$$\hat{\omega}_{t,t-h}^{\text{BG5}} = \frac{\hat{\Sigma}_{et,t-h}^{-1}}{t' \hat{\Sigma}_{et,t-h}^{-1} t}, \tag{51}$$

$$\hat{\Sigma}_{et,t-h}[i, j] = \sum_{\tau=h+1}^t \lambda^\tau e_{\tau,\tau-h,i} e_{\tau,\tau-h,j}.$$

Putting more weight on recent data means reducing the weight on past data and tends to increase the variance of the parameter estimates. Hence it will typically lead to poorer performance if the underlying data generating process is truly covariance stationary. Conversely, the underlying time-variations have to be quite strong to justify not using an expanding window. See Pesaran and Timmermann (2005) for further analysis of this point.

Diebold and Pauly (1987) embed these schemes in a general weighted least squares setup that chooses combination weights to minimize the weighted average of forecast errors from the combination. Let  $e_{t,t-h}^c = y_t - \omega' \hat{y}_{t,t-h}$  be the forecast error from the combination. Then one can minimize

$$\sum_{t=h+1}^T \sum_{\tau=h+1}^T \gamma_{t,\tau} e_{t,t-h}^c e_{\tau,\tau-h}^c, \tag{52}$$

or equivalently,  $\mathbf{e}' \mathbf{\Gamma} \mathbf{e}^c$ , where  $\mathbf{\Gamma}$  is a  $(T - h) \times (T - h)$  matrix with  $[t, \tau]$  element  $\gamma_{t,\tau}$  and  $\mathbf{e}^c$  is a  $(T - h) \times 1$  vector of errors from the forecast combination. Assuming that  $\mathbf{\Gamma}$  is diagonal, equal-weights on all past observations correspond to  $\gamma_{tt} = 1$  for all  $t$ , linearly declining weights can be represented as  $\gamma_{tt} = t$ , and geometrically declining weights take the form  $\gamma_{tt} = \lambda^{T-t}$ ,  $0 < \lambda \leq 1$ . Finally, Diebold and Pauly introduce two new weighting schemes, namely nonlinearly declining weights,  $\gamma_{tt} = t^\lambda$ ,  $\lambda \geq 0$ , and the Box–Cox transform weights

$$\gamma_{tt} = \begin{cases} (t^\lambda - 1)/\lambda & \text{if } 0 < \lambda \leq 1, \\ \ln(t) & \text{if } \lambda = 0. \end{cases}$$

These weights can be either declining at an increasing rate or at a decreasing rate, depending on the sign of  $\lambda - 1$ . This is clearly an attractive feature and one that, e.g., the geometrically declining weights do not have.

Diebold and Pauly also consider regression-based combinations with time-varying parameters. For example, if both the intercept and slope of the combination regression are allowed to vary over time,

$$\hat{y}_{t+h} = \sum_{i=1}^N (g_t^i + \mu_t^i) \hat{y}_{t+h,t,i},$$

where  $g^i(t) + \mu_t^i$  represent random variation in the combination weights. This approach explicitly models the evolution in the combination weights as opposed to doing this indirectly through the weighting of past and current forecast errors.

Instead of using adaptive schemes for updating the parameter estimates, an alternative is to explicitly model time-variations in the combination weights. A class of combination schemes considered by, e.g., Sessions and Chattererjee (1989), Zellner, Hong and Min (1991) and LeSage and Magura (1992) lets the combination weights evolve smoothly according to a time-varying parameter model:

$$\begin{aligned}
 y_{t+h} &= \tilde{\omega}'_{t+h,t} \mathbf{z}_{t+h} + \varepsilon_{t+h}, \\
 \tilde{\omega}_{t+h,t} &= \tilde{\omega}_{t,t-h} + \boldsymbol{\eta}_{t+h},
 \end{aligned}
 \tag{53}$$

where  $\mathbf{z}_{t+h} = (1 \hat{\mathbf{y}}'_{t+h,t})'$  and  $\tilde{\omega}_{t+h,t} = (\omega_{0t+h,t} \boldsymbol{\omega}'_{1t+h,t})'$ . It is typically assumed that (for  $h = 1$ )  $\varepsilon_{t+h} \sim \text{iid}(0, \sigma_\varepsilon^2)$ ,  $\boldsymbol{\eta}_{t+h} \sim \text{iid}(0, \boldsymbol{\Sigma}_\eta^2)$  and  $\text{Cov}(\varepsilon_{t+h}, \boldsymbol{\eta}_{t+h}) = \mathbf{0}$ .

Changes in the combination weights may instead occur more discretely, driven by some switching indicator,  $I_e$ , cf. Deutsch, Granger and Terasvirta (1994):

$$y_{t+h} = I_{\mathbf{e}_t \in A} (\omega_{01} + \boldsymbol{\omega}'_1 \hat{\mathbf{y}}_{t+h,t}) + (1 - I_{\mathbf{e}_t \in A}) (\omega_{02} + \boldsymbol{\omega}'_2 \hat{\mathbf{y}}_{t+h,t}) + \varepsilon_{t+h}.
 \tag{54}$$

Here  $\mathbf{e}_t = \boldsymbol{\iota} y_t - \hat{\mathbf{y}}_{t,t-h}$  is the vector of period- $t$  forecast errors;  $I_{\mathbf{e}_t \in A}$  is an indicator function taking the value unity when  $\mathbf{e}_t \in A$  and zero otherwise, for  $A$  some pre-defined set defining the switching condition. This provides a broad class of time-varying combination schemes as  $I_{\mathbf{e}_t \in A}$  can depend on past forecast errors or other variables in a number of ways. For example,  $I_{\mathbf{e}_t \in A}$  could be unity if the forecast error is positive, zero otherwise.

Engle, Granger and Kraft (1984) propose time-varying combining weights that follow a bivariate ARCH scheme and are constrained to sum to unity. They assume that the distribution of the two forecast errors  $\mathbf{e}_{t+h,t} = (e_{t+h,t,1} \ e_{t+h,t,2})'$  is bivariate Gaussian  $N(\mathbf{0}, \boldsymbol{\Sigma}_{t+h,t})$  where  $\boldsymbol{\Sigma}_{t+h,t}$  is the conditional covariance matrix.

A flexible mixture model for time-variation in the combination weights has been proposed by Elliott and Timmermann (2005). This approach is able to track both sudden and discrete as well as more gradual shifts in the joint distribution of  $(y_{t+h} \ \hat{\mathbf{y}}'_{t+h,t})$ . Suppose that the joint distribution of  $(y_{t+h} \ \hat{\mathbf{y}}'_{t+h,t})$  is driven by an unobserved state variable,  $S_{t+h}$ , which assumes one of  $n_s$  possible values, i.e.  $S_{t+h} \in (1, \dots, n_s)$ . Conditional on a given realization of the underlying state,  $S_{t+h} = s_{t+h}$ , the joint distribution of  $y_{t+h}$  and  $\hat{\mathbf{y}}_{t+h,t}$  is assumed to be Gaussian

$$\left. \begin{pmatrix} y_{t+h} \\ \hat{\mathbf{y}}_{t+h,t} \end{pmatrix} \right|_{S_{t+h}} \sim N \left( \begin{pmatrix} \mu_{yS_{t+h}} \\ \boldsymbol{\mu}_{\hat{\mathbf{y}}S_{t+h}} \end{pmatrix}, \begin{pmatrix} \sigma_{yS_{t+h}}^2 & \boldsymbol{\sigma}'_{y\hat{\mathbf{y}}S_{t+h}} \\ \boldsymbol{\sigma}_{y\hat{\mathbf{y}}S_{t+h}} & \boldsymbol{\Sigma}_{\hat{\mathbf{y}}\hat{\mathbf{y}}S_{t+h}} \end{pmatrix} \right).
 \tag{55}$$

This is similar to (7) but now conditional on  $S_{t+h}$ , which is important. This model generalizes (28) to allow for an arbitrary number of states. State transitions are assumed

to be driven by a first-order Markov chain  $\mathbf{P} = \Pr(S_{t+h} = s_{t+h} | S_t = s_t)$

$$\mathbf{P} = \begin{pmatrix} p_{11} & p_{12} & \cdots & p_{1n_s} \\ p_{21} & p_{22} & \cdots & \vdots \\ \vdots & \vdots & \cdots & p_{n_s-1n_s} \\ p_{n_s1} & \cdots & p_{n_s n_s-1} & p_{n_s n_s} \end{pmatrix}. \quad (56)$$

Conditional on  $S_{t+h} = s_{t+h}$ , the expectation of  $y_{t+h}$  is linear in the prediction signals,  $\hat{\mathbf{y}}_{t+h,t}$ , and thus takes the form of state-dependent intercept and combination weights:

$$\mathbb{E}[y_{t+h} | \hat{\mathbf{y}}_{t+h,t}, S_{t+h}] = \mu_{y s_{t+h}} + \boldsymbol{\sigma}'_{y \hat{\mathbf{y}}_{s_{t+h}}} \boldsymbol{\Sigma}_{\hat{\mathbf{y}}_{s_{t+h}}}^{-1} (\hat{\mathbf{y}}_{t+h,t} - \boldsymbol{\mu}_{\hat{\mathbf{y}}_{s_{t+h}}}). \quad (57)$$

Accounting for the fact that the underlying state is unobservable, the conditionally expected loss given current information,  $\mathcal{F}_t$ , and state probabilities,  $\pi_{s_{t+h},t}$ , becomes:

$$\mathbb{E}[e_{t+h}^2 | \pi_{s_{t+h},t}, \mathcal{F}_t] = \sum_{s_{t+h}=1}^{n_s} \pi_{s_{t+h},t} \{ \mu_{e s_{t+h}}^2 + \sigma_{e s_{t+h}}^2 \}, \quad (58)$$

where  $\pi_{s_{t+h},t} = \Pr(S_{t+h} = s_{t+h} | \mathcal{F}_t)$  is the probability of being in state  $s_{t+h}$  in period  $t+h$  conditional on current information,  $\mathcal{F}_t$ . Assuming a linear combination conditional on  $\mathcal{F}_t$ ,  $\pi_{s_{t+h},t}$  the optimal combination weights,  $\omega_{0t+h,t}^*$ ,  $\boldsymbol{\omega}_{t+h,t}^*$  become [cf. Elliott and Timmermann (2005)]

$$\begin{aligned} \omega_{0t+h,t}^* &= \sum_{s_{t+h}=1}^{n_s} \pi_{s_{t+h},t} \mu_{y s_{t+h}} - \left( \sum_{s_{t+h}=1}^{n_s} \pi_{s_{t+h},t} \boldsymbol{\mu}'_{\hat{\mathbf{y}}_{s_{t+h}}} \right) \boldsymbol{\omega}_{th} \equiv \bar{\mu}_{y t} - \bar{\boldsymbol{\mu}}'_{\hat{\mathbf{y}} t} \boldsymbol{\omega}_{th}, \\ \boldsymbol{\omega}_{t+h,t}^* &= \left( \sum_{s_{t+h}=1}^{n_s} \pi_{s_{t+h},t} (\boldsymbol{\mu}_{\hat{\mathbf{y}}_{s_{t+h}}} \boldsymbol{\mu}'_{\hat{\mathbf{y}}_{s_{t+h}}} + \boldsymbol{\Sigma}_{\hat{\mathbf{y}}_{s_{t+h}}}) - \bar{\boldsymbol{\mu}}_{\hat{\mathbf{y}} t} \bar{\boldsymbol{\mu}}'_{\hat{\mathbf{y}} t} \right)^{-1} \\ &\quad \times \left( \sum_{s_{t+h}=1}^{n_s} \pi_{s_{t+h},t} (\mu_{y s_{t+h}} \boldsymbol{\mu}_{\hat{\mathbf{y}}_{s_{t+h}}} + \boldsymbol{\sigma}_{y \hat{\mathbf{y}}_{s_{t+h}}}) - \bar{\mu}_{y t+h,t} \bar{\boldsymbol{\mu}}'_{\hat{\mathbf{y}} t+h,t} \right), \end{aligned} \quad (59)$$

where  $\bar{\mu}_{y t+h,t} = \sum_{s_{t+h}=1}^{n_s} \pi_{s_{t+h},t} \mu_{y s_{t+h}}$  and  $\bar{\boldsymbol{\mu}}_{\hat{\mathbf{y}} t+h,t} = \sum_{s_{t+h}=1}^{n_s} \pi_{s_{t+h},t} \boldsymbol{\mu}_{\hat{\mathbf{y}}_{s_{t+h}}}$ . The standard weights in (8) can readily be obtained by setting  $n_s = 1$ .

It follows from (59) that the (conditionally) optimal combination weights will vary as the state probabilities vary over time as a function of the arrival of new information provided that  $\mathbf{P}$  is of rank greater than one.

#### 4.2. Nonlinear combination schemes

Two types of nonlinearities can be considered in forecast combinations. First, nonlinear functions of the forecasts can be used in the combination which is nevertheless linear in the unknown parameters:

$$\hat{y}_{t+h,t}^c = \omega_0 + \boldsymbol{\omega}' C(\hat{\mathbf{y}}_{t+h,t}). \quad (60)$$

Here  $C(\hat{\mathbf{y}}_{t+h,t})$  is a function of the underlying forecasts that typically includes a lead term that is linear in  $\hat{\mathbf{y}}_{t+h,t}$  in addition to higher order terms similar to a Volterra or Taylor series expansion. The nonlinearity in (60) only enters through the shape of the transformation  $C(\cdot)$  so the unknown parameters can readily be estimated by OLS although the small-sample properties of such estimates could be an issue due to possible outliers. A second and more general combination method considers nonlinearities in the combination parameters, i.e.

$$\hat{y}_{t+h,t}^c = C(\hat{\mathbf{y}}_{t+h,t}, \boldsymbol{\omega}). \quad (61)$$

There does not appear to be much work in this area, possibly because estimation errors already appear to be large in linear combination schemes. They can be expected to be even larger for nonlinear combinations whose parameters are generally less robust and more sensitive to outliers than those of the linear schemes. Techniques from the Handbook Chapter 9 by White (2006) could be readily used in this context, however.

One paper that does estimate nonlinear combination weights is the study by Donaldson and Kamstra (1996). This uses artificial neural networks to combine volatility forecasts from a range of alternative models. Their combination scheme takes the form

$$\begin{aligned} \hat{y}_{t+h,t}^c &= \beta_0 + \sum_{j=1}^N \beta_j \hat{y}_{t+h,t,j} + \sum_{i=1}^p \delta_i g(\mathbf{z}_{t+h,t} \boldsymbol{\gamma}_i), \\ g(\mathbf{z}_{t+h,t} \boldsymbol{\gamma}_i) &= \left( 1 + \exp \left( - \left( \gamma_{0,i} + \sum_{j=1}^N \gamma_{1,j} z_{t+h,t,j} \right) \right) \right)^{-1}, \\ z_{t+h,t,j} &= (\hat{y}_{t+h,t,j} - \bar{y}_{t+h,t}) / \hat{\sigma}_{y_{t+h,t}}, \\ p &\in \{0, 1, 2, 3\}. \end{aligned} \quad (62)$$

Here  $\bar{y}_{t+h,t}$  is the sample estimate of the mean of  $y$  across the forecasting models while  $\hat{\sigma}_{y_{t+h,t}}$  is the sample estimate of the standard deviation using data up to time  $t$ . This network uses logistic nodes. The linear model is nested as a special case when  $p = 0$  so no nonlinear terms are included. In an out-of-sample forecasting experiment for volatility in daily stock returns, Donaldson and Kamstra find evidence that the neural net combination applied to two underlying forecasts (a moving average variance model and a GARCH(1,1) model) outperforms traditional combination methods.

## 5. Shrinkage methods

In cases where the number of forecasts,  $N$ , is large relative to the sample size,  $T$ , the sample covariance matrix underlying standard combinations is subject to considerable estimation uncertainty. Shrinkage methods aim to trade off bias in the combination weights against reduced parameter estimation error in estimates of the combination

weights. Intuition for how shrinkage works is well summarized by [Ledoit and Wolf \(2004, p. 2\)](#): “The crux of the method is that those estimated coefficients in the sample covariance matrix that are extremely high tend to contain a lot of positive error and therefore need to be pulled downwards to compensate for that. Similarly, we compensate for the negative error that tends to be embedded inside extremely low estimated coefficients by pulling them upwards.” This problem can partially be resolved by imposing more structure on the estimator in a way that reduces estimation error although the key question remains how much and which structure to impose. Shrinkage methods let the forecast combination weights depend on the sample size relative to the number of cross-sectional models to be combined.

[Diebold and Pauly \(1990\)](#) propose to shrink towards equal-weights. Consider the standard linear regression model underlying most forecast combinations and for simplicity drop the time and horizon subscripts:

$$\mathbf{y} = \hat{\mathbf{y}}\boldsymbol{\omega} + \boldsymbol{\varepsilon}, \quad \boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma^2\mathbf{I}), \quad (63)$$

where  $\mathbf{y}$  and  $\boldsymbol{\varepsilon}$  are  $T \times 1$  vectors,  $\hat{\mathbf{y}}$  is the  $T \times N$  matrix of forecasts and  $\boldsymbol{\omega}$  is the  $N \times 1$  vector of combination weights. The standard normal-gamma conjugate prior  $\sigma^2 \sim IG(s_0^2, v_0)$ ,  $\boldsymbol{\omega}|\sigma \sim N(\boldsymbol{\omega}_0, \mathbf{M})$  implies that

$$P(\boldsymbol{\omega}, \sigma) \propto \sigma^{-N-v_0-1} \exp\left(\frac{-(v_0s_0^2 + (\boldsymbol{\omega} - \boldsymbol{\omega}_0)'\mathbf{M}(\boldsymbol{\omega} - \boldsymbol{\omega}_0))}{2\sigma^2}\right). \quad (64)$$

Under normality of  $\boldsymbol{\varepsilon}$  the likelihood function for the data is

$$L(\boldsymbol{\omega}, \sigma | \mathbf{y}, \hat{\mathbf{y}}) \propto \sigma^{-T} \exp\left(\frac{-(\mathbf{y} - \hat{\mathbf{y}}\boldsymbol{\omega})'(\mathbf{y} - \hat{\mathbf{y}}\boldsymbol{\omega})}{2\sigma^2}\right) \quad (65)$$

These results can be combined to give the marginal posterior for  $\boldsymbol{\omega}$  with mean

$$\bar{\boldsymbol{\omega}} = (\mathbf{M} + \hat{\mathbf{y}}'\hat{\mathbf{y}})^{-1}(\mathbf{M}\boldsymbol{\omega}_0 + \hat{\mathbf{y}}'\hat{\mathbf{y}}\hat{\boldsymbol{\omega}}), \quad (66)$$

where  $\hat{\boldsymbol{\omega}} = (\hat{\mathbf{y}}'\hat{\mathbf{y}})^{-1}\hat{\mathbf{y}}'\mathbf{y}$  is the least squares estimate of  $\boldsymbol{\omega}$ . Using a prior for  $\mathbf{M}$  that is proportional to  $\hat{\mathbf{y}}'\hat{\mathbf{y}}$ ,  $\mathbf{M} = g\hat{\mathbf{y}}'\hat{\mathbf{y}}$ , we get

$$\bar{\boldsymbol{\omega}} = (g\hat{\mathbf{y}}'\hat{\mathbf{y}} + \hat{\mathbf{y}}'\hat{\mathbf{y}})^{-1}(g\hat{\mathbf{y}}'\hat{\mathbf{y}}\boldsymbol{\omega}_0 + \hat{\mathbf{y}}'\hat{\mathbf{y}}\hat{\boldsymbol{\omega}}),$$

which can be used to obtain

$$\bar{\boldsymbol{\omega}} = \boldsymbol{\omega}_0 + \frac{\hat{\boldsymbol{\omega}} - \boldsymbol{\omega}_0}{1 + g}. \quad (67)$$

Clearly, the larger the value of  $g$ , the stronger the shrinkage towards the mean of the prior,  $\boldsymbol{\omega}_0$ , whereas small values of  $g$  suggest putting more weight on the data.

Alternatively, empirical Bayes methods can be used to estimate  $g$ . Suppose the prior for  $\boldsymbol{\omega}$  conditional on  $\sigma$  is Gaussian  $N(\boldsymbol{\omega}_0, \tau^2\mathbf{A}^{-1})$ . Then the posterior for  $\boldsymbol{\omega}$  is also Gaussian,  $N(\bar{\boldsymbol{\omega}}, \boldsymbol{\tau}^{-2}\mathbf{A} + \sigma^{-2}\hat{\mathbf{y}}'\hat{\mathbf{y}})$  and  $\sigma^2$  and  $\tau^2$  can be replaced by the estimates [cf.

Diebold and Pauly (1990)]

$$\hat{\sigma}^2 = \frac{(\mathbf{y} - \hat{\mathbf{y}}\hat{\boldsymbol{\omega}})'(\mathbf{y} - \hat{\mathbf{y}}\hat{\boldsymbol{\omega}})}{T},$$

$$\hat{\tau}^2 = \frac{(\hat{\boldsymbol{\omega}} - \boldsymbol{\omega}_0)'(\hat{\boldsymbol{\omega}} - \boldsymbol{\omega}_0)}{\text{tr}(\hat{\mathbf{y}}'\hat{\mathbf{y}})^{-1}} - \hat{\sigma}^2.$$

This gives rise to an empirical Bayes estimator of  $\boldsymbol{\omega}$  whose posterior mean is

$$\bar{\boldsymbol{\omega}} = \boldsymbol{\omega}_0 + \left( \frac{\hat{\tau}^2}{\hat{\sigma}^2 + \hat{\tau}^2} \right) (\hat{\boldsymbol{\omega}} - \boldsymbol{\omega}_0). \quad (68)$$

The empirical Bayes combination shrinks  $\hat{\boldsymbol{\omega}}$  towards  $\boldsymbol{\omega}_0$  and amounts to setting  $g = \hat{\sigma}^2/\hat{\tau}^2$  in (67). Notice that if  $\hat{\sigma}^2/\hat{\tau}^2 \rightarrow 0$ , the OLS estimator is obtained while if  $\hat{\sigma}^2/\hat{\tau}^2 \rightarrow \infty$ , the prior estimate  $\boldsymbol{\omega}_0$  is obtained as a special case. Diebold and Pauly argue that the combination weights should be shrunk towards the equal-weighted (simple) average so the combination procedure gives a convex combination of the least-squares and equal weights.

Stock and Watson (2004) also propose shrinkage towards the arithmetic average of forecasts. Let  $\hat{\omega}_{T,T-h,i}$  be the least-squares estimator of the weight on the  $i$ th model in the forecast combination based on data up to period  $T$ . The combination weights considered by Stock and Watson take the form (assuming  $T > h + N + 1$ )

$$\omega_{T,T-h,i} = \psi \hat{\omega}_{T,T-h,i} + (1 - \psi)(1/N),$$

$$\psi = \max(0, 1 - \kappa N / (T - h - N - 1)),$$

where  $\kappa$  regulates the strength of the shrinkage. Stock and Watson consider values  $\kappa = 1/4, 1/2$  or  $1$ . As the sample size,  $T$ , rises relative to  $N$ , the least squares estimate gets a larger weight. Indeed, if  $T$  grows at a faster rate than  $N$ , the least squares weight will, in the limit, get a weight of unity.

### 5.1. Shrinkage and factor structure

In a portfolio application Ledit and Wolf (2003) propose to shrink the weights towards a point implied by a single factor structure common from finance.<sup>12</sup> Suppose that the

<sup>12</sup> The problem of forming mean–variance efficient portfolios in finance is mathematically equivalent to that of combining forecasts, cf. Dunis, Timmermann and Moody (2001). In finance, the standard optimization problem minimizes the portfolio variance  $\boldsymbol{\omega}'\boldsymbol{\Sigma}\boldsymbol{\omega}$  subject to a given portfolio return,  $\boldsymbol{\omega}'\boldsymbol{\mu} = \mu_0$ , where  $\boldsymbol{\mu}$  is a vector of mean returns while  $\boldsymbol{\Sigma}$  is the covariance matrix of asset returns. Imposing also the constraint that the portfolio weights sum to unity, we have

$$\begin{aligned} \min_{\boldsymbol{\omega}} \quad & \boldsymbol{\omega}'\boldsymbol{\Sigma}\boldsymbol{\omega} \\ \text{s.t.} \quad & \boldsymbol{\omega}'\mathbf{1} = 1, \\ & \boldsymbol{\omega}'\boldsymbol{\mu} = \mu_0. \end{aligned}$$

individual forecast errors are affected by a single common factor,  $f_{et}$ :

$$e_{it} = \alpha_i + \beta_i f_{et} + \varepsilon_{it} \quad (69)$$

where the idiosyncratic residuals,  $\varepsilon_{it}$ , are assumed to be orthogonal across forecasting models and uncorrelated with  $f_{et}$ . This single factor model has a long tradition in finance but is also a natural starting point for forecasting purposes since forecast errors are generally strongly positively correlated. Letting  $\sigma_{f_e}^2$  be the variance of  $f_{et}$ , the covariance matrix of the forecast errors becomes

$$\Sigma_{ef} = \sigma_{f_e}^2 \boldsymbol{\beta} \boldsymbol{\beta}' + \mathbf{D}_\varepsilon, \quad (70)$$

where  $\boldsymbol{\beta} = (\beta_1 \dots \beta_N)'$  is the vector of factor sensitivities, while  $\mathbf{D}_\varepsilon$  is a diagonal matrix with the individual values of  $\text{Var}(\varepsilon_{it})$  on the diagonal. Estimation of  $\Sigma_{ef}$  requires determining only  $2N + 1$  parameters. Consistent estimates of these parameters are easily obtained by estimating (69) by OLS, equation by equation, to get

$$\hat{\Sigma}_{ef} = \hat{\sigma}_{f_e}^2 \hat{\boldsymbol{\beta}} \hat{\boldsymbol{\beta}}' + \hat{\mathbf{D}}_\varepsilon.$$

Typically this covariance matrix is biased due to the assumption that  $\mathbf{D}_\varepsilon$  is diagonal. For example, there may be more than a single common factor in the forecast errors and some forecasts may omit the same relevant variable in which case blocks of forecast errors will be correlated. Though biased, the single factor covariance matrix is typically surrounded by considerably smaller estimation errors than the unconstrained matrix,  $E[\mathbf{e}\mathbf{e}']$ , which can be estimated by

$$\hat{\Sigma}_e = \frac{1}{T-h} \sum_{\tau=h}^T \mathbf{e}_{\tau, \tau-h} \mathbf{e}'_{\tau, \tau-h},$$

where  $\mathbf{e}_{\tau, \tau-h}$  is an  $N \times 1$  vector of forecast errors. This estimator requires estimating  $N(N+1)/2$  parameters. Using  $\hat{\Sigma}_{ef}$  as the shrinkage point, Ledoit and Wolf (2003) propose minimizing the following quadratic loss as a function of the shrinkage parameter,  $\alpha$ ,

$$L(\alpha) = \|\alpha \hat{\Sigma}_{ef} + (1-\alpha) \hat{\Sigma}_e - \Sigma_e\|^2,$$

where  $\|\cdot\|^2$  is the Frobenius norm, i.e.  $\|\mathbf{Z}\|^2 = \text{trace}(\mathbf{Z}^2)$ ,  $\hat{\Sigma}_e = (1/T)\mathbf{e}(\mathbf{I} - \mathbf{u}'/T)\mathbf{e}'$  is the sample covariance matrix and  $\Sigma_e$  is the true matrix of squared forecast errors,  $E[\mathbf{e}\mathbf{e}']$ ,

This problem has the solution

$$\boldsymbol{\omega}^* = \boldsymbol{\Sigma}^{-1}(\boldsymbol{\mu})[(\boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\boldsymbol{\mu})]^{-1} \begin{pmatrix} \mu_0 \\ 1 \end{pmatrix}.$$

In the forecast combination problem the constraint  $\boldsymbol{\omega}'\mathbf{1} = 1$  is generally interpreted as guaranteeing an unbiased combined forecast – assuming of course that the individual forecasts are also unbiased. The only difference to the optimal solution from the forecast combination problem is that a minimum variance portfolio is derived for each separate value of the mean portfolio return,  $\mu_0$ .

where  $\mathbf{e}$  is an  $N \times T$  matrix of forecast errors. Letting  $\hat{f}_{ij}$  be the  $(i, j)$  entry of  $\hat{\Sigma}_{ef}$ ,  $\hat{\sigma}_{ij}$  the  $(i, j)$  element of  $\hat{\Sigma}_e$  and  $\phi_{ij}$  the  $(i, j)$  element of the single factor covariance matrix,  $\Sigma_{ef}$ , while  $\sigma_{ij}$  is the  $(i, j)$  element of  $\Sigma_e$ , they demonstrate that the optimal shrinkage takes the form

$$\alpha^* = \frac{1}{T} \frac{\pi - \rho}{\gamma} + O\left(\frac{1}{T^2}\right),$$

where

$$\begin{aligned} \pi &= \sum_{i=1}^N \sum_{j=1}^N \text{AsyVar}(\sqrt{T} \hat{\sigma}_{ij}), \\ \rho &= \sum_{i=1}^N \sum_{j=1}^N \text{AsyCov}(\sqrt{T} \hat{f}_{ij}, \sqrt{T} \hat{\sigma}_{ij}), \\ \gamma &= \sum_{i=1}^N \sum_{j=1}^N (\phi_{ij} - \sigma_{ij})^2. \end{aligned}$$

Hence,  $\pi$  measures the (scaled) sum of asymptotic variances of the sample covariance matrix ( $\hat{\Sigma}_e$ ),  $\rho$  measures the (scaled) sum of asymptotic covariances of the sample covariance matrix ( $\hat{\Sigma}_e$ ) and the single-factor covariance matrix ( $\hat{\Sigma}_{ef}$ ), while  $\gamma$  measures the degree of misspecification (bias) in the single factor model. Ledoit and Wolf propose consistent estimators  $\hat{\pi}$ ,  $\hat{\rho}$  and  $\hat{\gamma}$  under the assumption of IID forecast errors.<sup>13</sup>

## 5.2. Constraints on combination weights

Shrinkage bears an interesting relationship to portfolio weight constraints in finance. It is commonplace to consider minimization of portfolio variance subject to a set of equality and inequality constraints on the portfolio weights. Portfolio weights are often constrained to be non-negative (due to no short selling) and not to exceed certain upper bounds (due to limits on ownership in individual stocks). Reflecting this, let  $\hat{\Sigma}$  be an estimate of the covariance matrix for some cross-section of asset returns with row  $i$ , column  $j$  element  $\hat{\Sigma}[i, j]$  and consider the optimization program

$$\begin{aligned} \boldsymbol{\omega}^* &= \arg \min_{\boldsymbol{\omega}} \frac{1}{2} \boldsymbol{\omega}' \hat{\Sigma} \boldsymbol{\omega} \\ \text{s.t. } \boldsymbol{\omega}' \mathbf{1} &= 1, \\ \omega_i &\geq 0, \quad i = 1, \dots, N, \\ \omega_i &\leq \bar{\omega}, \quad i = 1, \dots, N. \end{aligned} \tag{71}$$

<sup>13</sup> It is worth pointing out that the assumption that  $\mathbf{e}$  is IID is unlikely to hold for forecast errors which could share common dynamics in first, second or higher order moments or even be serially correlated, cf. Diebold (1988).



This gives a set of Kuhn–Tucker conditions:

$$\sum_j \hat{\Sigma}[i, j] \omega_j - \lambda_i + \delta_i = \lambda_0 \geq 0, \quad i = 1, \dots, N,$$

$$\lambda_i \geq 0 \quad \text{and} \quad \lambda_i = 0 \quad \text{if} \quad \omega_i > 0,$$

$$\delta_i \geq 0 \quad \text{and} \quad \delta_i = 0 \quad \text{if} \quad \omega_i < \bar{\omega}.$$

Lagrange multipliers for the lower and upper bounds are collected in the vectors  $\lambda = (\lambda_1, \dots, \lambda_N)'$  and  $\delta = (\delta_1, \dots, \delta_N)'$ ;  $\lambda_0$  is the Lagrange multiplier for the constraint that the weights sum to one.

Constraints on combination weights effectively have two effects. First, they shrink the largest elements of the covariance matrix towards zero. This reduces the effects of estimation error that can be expected to be strongest for assets with extreme weights. The second effect is that it may introduce specification errors to the extent that the true population values of the optimal weights actually lie outside the assumed interval.

Jaganathan and Ma (2003) show the following result. Let

$$\tilde{\Sigma} = \hat{\Sigma} + (\delta \iota' + \iota \delta') - (\lambda \iota' + \iota \lambda'). \quad (72)$$

Then  $\tilde{\Sigma}$  is symmetric and positive semi-definite. Constructing a solution to the inequality constrained problem (71) is shown to be equivalent to finding the optimal weights for the unconstrained quadratic form based on the modified covariance matrix in (72)  $\tilde{\Sigma} = \hat{\Sigma} + (\delta \iota' + \iota \delta') - (\lambda \iota' + \iota \lambda')$ .

Furthermore, it turns out that  $\tilde{\Sigma}$  can be interpreted as a shrinkage version of  $\hat{\Sigma}$ . To see this, consider the weights that are affected by the lower bound so  $\tilde{\Sigma} = \hat{\Sigma} - (\lambda \iota' + \iota \lambda')$ . When the constraint for the lower bound is binding (so a combination weight would have been negative), the covariances of a particular forecast error with all other errors are reduced by the strictly positive Lagrange multipliers and its variance is shrunk. Imposing the non-negativity constraints shrinks the largest covariance estimates that would have resulted in negative weights. Since the largest estimates of the covariance are more likely to be the result of estimation error, such shrinkage can have the effect of reducing estimation error and have the potential to improve out-of-sample performance of the combination.

In the case of the upper bounds, those forecasts whose unconstrained weights would have exceeded  $\bar{\omega}$  are also the ones for which the variance and covariance estimates tend to be smallest. These forecasts have strictly positive Lagrange multipliers on the upper bound constraint, meaning that their forecast error variance will be increased by  $2\delta_i$  while the covariances in the modified covariance matrix  $\tilde{\Sigma}$  will be increased by  $\delta_i + \delta_j$ . Again this corresponds to shrinkage towards the cross-sectional average of the variances and covariances.

## 6. Combination of interval and probability distribution forecasts

So far we have focused on combining point forecasts. This, of course, reflects the fact that the vast majority of academic studies on forecasting only report point forecasts. However, there has been a growing interest in studying interval and probability distribution forecasts and an emerging literature in economics is considering the scope for using combination methods for such forecasts. This is preceded by the use of combined probability forecasting in areas such as meteorology, cf. Sanders (1963). Genest and Zidek (1986) present a broad survey of various techniques in this area.

### 6.1. The combination decision

As in the case of combinations of point forecasts it is natural to ask whether the best strategy is to use only a single probability forecast or a combination of these. This is related to the concept of forecast encompassing which generalizes from point to density forecasts as follows. Suppose we are considering combining  $N$  distribution forecasts  $f_1, \dots, f_N$  whose joint distribution with  $y$  is  $P(y, f_1, f_2, \dots, f_N)$ . Factoring this into the product of the conditional distribution of  $y$  given  $f_1, \dots, f_N$ ,  $P(y|f_1, \dots, f_N)$ , and the marginal distribution of the forecasts,  $P(f_1, \dots, f_N)$ , we have

$$P(y, f_1, f_2, \dots, f_N) = P(y|f_1, \dots, f_N)P(f_1, \dots, f_N). \quad (73)$$

A probability forecast that does not provide information about  $y$  given all the other probability density forecasts is referred to as extraneous by Clemen, Murphy and Winkler (1995). If the  $i$ th forecast is extraneous we must have

$$P(y|f_1, f_2, \dots, f_N) = P(y|f_1, f_2, \dots, f_{i-1}, f_{i+1}, \dots, f_N). \quad (74)$$

If (74) holds, probability forecast  $f_i$  does not contain any information that is useful for forecasting  $y$  given the other  $N - 1$  probability forecasts. Only if forecast  $i$  does not satisfy (74) does it follow that this model is not encompassed by the other models. Interestingly, adding more forecasting models (i.e. increasing  $N$ ) can lead a previously extraneous model to become non-extraneous if it contains information about the relationship between the existing  $N - 1$  methods and the new forecasts.

For pairwise comparison of probability forecasts, Clemen, Murphy and Winkler (1995) define the concept of sufficiency. This concept is important because if forecast 1 is sufficient for forecast 2, then 1's forecasts will be of greater value to *all* users than forecast 2. Conversely, if neither model is sufficient for the other we would expect some forecast users to prefer model 1 while others prefer model 2. To illustrate this concept, consider two probability forecasts,  $f_1 = P_1(x = 1)$  and  $f_2 = P_2(x = 1)$  of some event,  $X$ , where  $x = 1$  if the event occurs while it is zero otherwise. Also let  $v_1(f) = P(f_1 = f)$  and  $v_2(g) = P(f_2 = g)$ , where  $f, g \in \mathcal{G}$ , and  $\mathcal{G}$  is the set of permissible probabilities. Forecast 1 is then said to be sufficient for forecast 2 if there

exists a stochastic transformation  $\zeta(g|f)$  such that for all  $g \in \mathcal{G}$ ,

$$\begin{aligned}\sum_f \zeta(g|f)v_1(f) &= v_2(g), \\ \sum_f \zeta(g|f)fv_1(f) &= gv_2(g).\end{aligned}$$

The function  $\zeta(g|f)$  is said to be a stochastic transformation provided that it lies between zero and one and integrates to unity. It represents an additional randomization that has the effect of introducing noise into the first forecast.

## 6.2. Combinations of probability density forecasts

Combinations of probability density or distribution forecasts impose new requirements beyond those we saw for combinations of point forecasts, namely that the combination must be convex with weights confined to the zero-one interval so that the probability forecast never becomes negative and always sums to one.

This still leaves open a wide set of possible combination schemes. An obvious way to combine a collection of probability forecasts  $\{F_{t+h,t,1}, \dots, F_{t+h,t,N}\}$  is through the convex combination (“linear opinion pool”):

$$\bar{F}^c = \sum_{i=1}^N \omega_{t+h,t,i} F_{t+h,t,i}, \quad (75)$$

with  $0 \leq \omega_{t+h,t,i} \leq 1$  ( $i = 1, \dots, N$ ) and  $\sum_{i=1}^N \omega_{t+h,t,i} = 1$  to ensure that the combined probability forecast is everywhere non-negative and integrates to one. The generalized linear opinion pool adds an extra probability forecast,  $F_{t+h,t,0}$ , and takes the form

$$\bar{F}^c = \sum_{i=0}^N \omega_{t+h,t,i} F_{t+h,t,i}. \quad (76)$$

Under this scheme the weights are allowed to be negative  $\omega_0, \omega_1, \dots, \omega_n \in [-1, 1]$  although they still are restricted to sum to unity:  $\sum_{i=0}^N \omega_{t+h,t,i} = 1$ .  $F_{t+h,t,0}$  can be shown to exist under conditions discussed by [Genest and Zidek \(1986\)](#).

Alternatively, one can adopt a logarithmic combination of densities

$$\bar{f}^l = \prod_{i=1}^N f_{t+h,t,i}^{\omega_{t+h,t,i}} \bigg/ \int \prod_{i=1}^N f_{t+h,t,i}^{\omega_{t+h,t,i}} d\mu, \quad (77)$$

where  $\{\omega_{t+h,t,1}, \dots, \omega_{t+h,t,N}\}$  are weights chosen such that the integral in the denominator is finite and  $\mu$  is the underlying probability measure. This combination is less dispersed than the linear combination and is also unimodal, cf. [Genest and Zidek \(1986\)](#).

### 6.3. Bayesian methods

Bayesian approaches have been widely used to construct combinations of probability forecasts. For example, [Min and Zellner \(1993\)](#) propose combinations based on posterior odds ratios. Let  $p_1$  and  $p_2$  be the posterior probabilities of two models (a fixed parameter and a time-varying parameter model in their application) while  $k = p_1/p_2$  is the posterior odds ratio of the two models. Assuming that the two models,  $M_1$  and  $M_2$ , are exhaustive the proposed combination scheme has a conditional mean of

$$\begin{aligned} E[y] &= p_1 E[y|M_1] + (1 - p_1) E[y|M_2] \\ &= \frac{k}{1+k} E[y|M_1] + \frac{1}{1+k} E[y|M_2]. \end{aligned} \quad (78)$$

[Palm and Zellner \(1992\)](#) propose a combination method that accounts for the full correlation structure between the forecast errors. They model the one-step forecast errors from the individual models as follows:

$$y_{t+1} - \hat{y}_{it+1,t} = \theta_i + \varepsilon_{it+1} + \eta_{t+1}, \quad (79)$$

where  $\theta_i$  is the bias in the  $i$ th model's forecast – reflecting perhaps the forecaster's asymmetric loss, cf. [Zellner \(1986\)](#) –  $\varepsilon_{it+1}$  is an idiosyncratic forecast error and  $\eta_{t+1}$  is a common component in the forecast errors reflecting an unpredictable component of the outcome variable. It is assumed that both  $\varepsilon_{it+1} \sim N(0, \sigma_\varepsilon^2)$  and  $\eta_{t+1} \sim N(0, \sigma_\eta^2)$  are serially uncorrelated (as well as mutually uncorrelated) Gaussian variables with zero mean.

For the case with zero bias ( $\theta_i = 0$ ), [Winkler \(1981\)](#) shows that when  $\varepsilon_{it+1} + \eta_{t+1}$  ( $i = 1, \dots, N$ ) has known covariance matrix,  $\Sigma_0$ , the predictive density function of  $y_{t+1}$  given an  $N$ -vector of forecasts  $\hat{\mathbf{y}}_{t+1,t} = (\hat{y}_{t+1,t,1}, \dots, \hat{y}_{t+1,t,N})'$  is Gaussian with mean  $\boldsymbol{\iota}' \Sigma_0^{-1} \hat{\mathbf{y}}_{t+1,t} / \boldsymbol{\iota}' \Sigma_0^{-1} \boldsymbol{\iota}$  and variance  $\boldsymbol{\iota}' \Sigma_0^{-1} \boldsymbol{\iota}$ . When the covariance matrix of the  $N$  time-varying parts of the forecast errors  $\varepsilon_{it+1} + \eta_{t+1}$ ,  $\Sigma$ , is unknown but has an inverted Wishart prior  $IW(\Sigma | \Sigma_0, \delta_0, N)$  with  $\delta_0 \geq N$ , the predictive distribution of  $y_{T+1}$  given  $\mathcal{F}_T = \{y_1, \dots, y_T, \hat{\mathbf{y}}_{2,1}, \dots, \hat{\mathbf{y}}_{T,T-1}, \hat{\mathbf{y}}_{T+1,T}\}$  is a univariate student-t with degrees of freedom parameter  $\delta_0 + N - 1$ , mean  $m^* = \boldsymbol{\iota}' \Sigma_0^{-1} \hat{\mathbf{y}}_{T+1,T} / \boldsymbol{\iota}' \Sigma_0^{-1} \boldsymbol{\iota}$  and variance  $(\delta_0 + N - 1) s^{*2} / (\delta_0 + N - 3)$ , where  $s^{*2} = (\delta_0 + (m^* \boldsymbol{\iota} - \hat{\mathbf{y}}_{T+1,T})' \Sigma_0^{-1} (m^* \boldsymbol{\iota} - \hat{\mathbf{y}}_{T+1,T})) / (\delta_0 + N - 1) \boldsymbol{\iota}' \Sigma_0^{-1} \boldsymbol{\iota}$ .

[Palm and Zellner \(1992\)](#) extend these results to allow for a non-zero bias. Given a set of  $N$  forecasts  $\hat{\mathbf{y}}_{t+1,t}$  over  $T$  periods they express the forecast errors  $y_t - \hat{y}_{t-1,t,i} = \theta_i + \varepsilon_{it} + \eta_t$  as a  $T \times N$  multivariate regression model:

$$\mathbf{Y} = \boldsymbol{\iota} \boldsymbol{\theta} + \mathbf{U}.$$

Suppose that the structure of the forecast errors (79) is reflected in a Wishart prior for  $\Sigma^{-1}$  with  $\nu$  degrees of freedom and covariance matrix  $\Sigma_0 = \Sigma_{\varepsilon_0} + \sigma_{\eta_0}^2 \boldsymbol{\iota} \boldsymbol{\iota}'$  (with known

parameters  $v, \Sigma_{\varepsilon 0}, \sigma_{\eta 0}^2$ ):

$$P(\Sigma^{-1}) \propto |\Sigma^{-1}|^{(v-N-1)/2} |\Sigma_0^{-1}|^{-v/2} \exp\left(-\frac{1}{2}\text{tr}(\Sigma_0 \Sigma^{-1})\right).$$

Assuming a sample of  $T$  observations and a likelihood function

$$L(\theta, \Sigma^{-1} | \mathcal{F}_T) \propto |\Sigma^{-1}|^{-T/2} \exp\left(-\frac{1}{2}\text{tr}(\mathbf{S} \Sigma^{-1}) - \frac{1}{2}\text{tr}((\theta - \hat{\theta}) \iota' (\theta - \hat{\theta})' \Sigma^{-1})\right),$$

where  $\hat{\theta} = (\iota' \iota)^{-1} \iota' \mathbf{Y}$  and  $\mathbf{S} = (\mathbf{Y} - \iota \hat{\theta}')' (\mathbf{Y} - \iota \hat{\theta}')$ , Palm and Zellner derive the predictive distribution function of  $y_{T+1}$  given  $\mathcal{F}_T$ :

$$P(y_{T+1} | \mathcal{F}_T) \propto [1 + (y_{T+1} - \bar{\mu})^2 / (T-1) s^{**2}]^{-(T+v)/2},$$

where  $\bar{\mu} = \iota' \bar{\mathbf{S}}^{-1} \hat{\boldsymbol{\mu}} / \iota' \bar{\mathbf{S}}^{-1} \iota$ ,  $s^{**2} = [T+1 + T(\bar{\mu} \iota - \hat{\boldsymbol{\mu}})' \bar{\mathbf{S}}^{-1} (\bar{\mu} \iota - \hat{\boldsymbol{\mu}})] / (T(T-1) \iota' \bar{\mathbf{S}}^{-1} \iota)$ ,  $\hat{\boldsymbol{\mu}} = \hat{y}_{T+1} - \hat{\theta}$  and  $\bar{\mathbf{S}} = \mathbf{S} + \Sigma_0$ . This approach provides a complete solution to the forecast combination problem that accounts for the joint distribution of forecast errors from the individual models.

### 6.3.1. Bayesian model averaging

Bayesian Model Averaging methods have been proposed by, inter alia, Leamer (1978), Raftery, Madigan and Hoeting (1997) and Hoeting et al. (1999) and are increasingly used in empirical studies, see, e.g., Jackson and Karlsson (2004). Under this approach, the predictive density can be computed by averaging over a set of models,  $i = 1, \dots, N$ , each characterized by parameters  $\theta_i$ :

$$f(y_{t+h} | \mathcal{F}_t) = \sum_{i=1}^N \Pr(M_i | \mathcal{F}_t) f_i(y_{t+h}, \theta_i | \mathcal{F}_t), \quad (80)$$

where  $\Pr(M_i | \mathcal{F}_t)$  is the posterior probability of model  $M_i$  obtained from the model priors  $\Pr(M_i)$ , the priors for the unknown parameters,  $\Pr(\theta_i | M_i)$ , and the likelihood functions of the models under consideration.  $f_i(y_{t+h}, \theta_i | \mathcal{F}_t)$  is the density of  $y_{t+h}$  and  $\theta_i$  under the  $i$ th model, given information at time  $t$ ,  $\mathcal{F}_t$ . Note that unlike the combination weights used for point forecasts such as (12), these weights do not account for correlations between forecasts. However, the approach is quite general and does not require the use of conjugate families of distributions. More details are provided in the Handbook Chapter 1 by Geweke and Whitemann (2006).

### 6.4. Combinations of quantile forecasts

Combinations of quantile forecasts do not pose any new issues except for the fact that the associated loss function used to combine quantiles is typically no longer continuous and differentiable. Instead predictions of the  $\alpha$ th quantile can be related to the ‘tick’ loss

function

$$L_\alpha(e_{t+h,t}) = (\alpha - 1_{e_{t+h,t} < 0})e_{t+h,t},$$

where  $1_{e_{t+h,t} < 0}$  is an indicator function taking a value of unity if  $e_{t+h,t} < 0$ , and is otherwise zero, cf. [Giacomini and Komunjer \(2005\)](#). Given a set of quantile forecasts  $q_{t+h,t,1}, \dots, q_{t+h,t,N}$ , quantile forecast combinations can then be based on formulas such as

$$q_{t+h,t}^c = \sum_{i=1}^N \omega_i q_{t+h,t,i},$$

possibly subject to constraints such as  $\sum_{i=1}^N \omega_i = 1$ .

More caution should be exercised when forming combinations of interval forecasts. Suppose that we have  $N$  interval forecasts each taking the form of a lower and an upper limit  $\{l_{t+h,t,i}; u_{t+h,t,i}\}$ . While weighted averages  $\{\bar{l}_{t+h,t,i}^c; \bar{u}_{t+h,t,i}^c\}$

$$\begin{aligned} \bar{l}_{t+h,t,i}^c &= \sum_{i=1}^N \omega_{t+h,t,i}^l l_{t+h,t,i}, \\ \bar{u}_{t+h,t,i}^c &= \sum_{i=1}^N \omega_{t+h,t,i}^u u_{t+h,t,i} \end{aligned} \tag{81}$$

may seem natural, they are not guaranteed to provide correct coverage rates. To see this, consider the following two 97% confidence intervals for a normal mean:

$$\begin{aligned} &\left[ \bar{y} - 2.58 \frac{\sigma}{T}, \bar{y} + 1.96 \frac{\sigma}{T} \right], \\ &\left[ \bar{y} - 1.96 \frac{\sigma}{T}, \bar{y} + 2.58 \frac{\sigma}{T} \right]. \end{aligned}$$

The average of these confidence intervals,  $[\bar{y} - 2.27 \frac{\sigma}{T}, \bar{y} + 2.27 \frac{\sigma}{T}]$  has a coverage of 97.7%. Combining confidence intervals may thus change the coverage rate.<sup>14</sup> The problem here is that the underlying end-points for the two forecasts (i.e.  $\bar{y} - 2.58 \frac{\sigma}{T}$  and  $\bar{y} - 1.96 \frac{\sigma}{T}$ ) are not estimates of the same quantiles. While it is natural to combine estimates of the same  $\alpha$ -quantile, it is less obvious that combination of forecast intervals makes much sense unless one can be assured that the end-points are lined up and are estimates of the same quantiles.

<sup>14</sup> I am grateful to Mark Watson for suggesting this example.

## 7. Empirical evidence

The empirical literature on forecast combinations is voluminous and includes work in several areas such as management science, economics, operations research, meteorology, psychology and finance. The work in economics dates back to Reid (1968) and Bates and Granger (1969). Although details and results vary across studies, it is possible to extract some broad conclusions from much of this work. Such conclusions come with a stronger than usual caveat *emtor* since for each point it is possible to construct counter examples. This is necessarily the case since findings depend on the number of models,  $N$  (as well as their type), the sample size,  $T$ , the extent of instability in the underlying data set and the structure of the covariance matrix of the forecast errors (e.g., diagonal or with similar correlations).

Nevertheless, empirical findings in the literature on forecast combinations broadly suggest that (i) simple combination schemes are difficult to beat. This is often explained by the importance of parameter estimation error in the combination weights. Consequently, methods aimed at reducing such errors (such as shrinkage or combination methods that ignore correlations between forecasts) tend to perform well; (ii) forecasts based exclusively on the model with the best in-sample performance often leads to poor out-of-sample forecasting performance; (iii) trimming of the worst models and clustering of models with similar forecasting performance prior to combination can yield considerable improvements in forecasting performance, especially in situations involving large numbers of forecasts; (iv) shrinkage to simple forecast combination weights often improves performance; and (v) some time-variation or adaptive adjustment in the combination weights (or perhaps in the underlying models being combined) can often improve forecasting performance. In the following we discuss each of these points in more detail. The section finishes with a brief empirical application to a large macroeconomic data set from the G7 economies.

### 7.1. Simple combination schemes are hard to beat

It has often been found that simple combinations – that is, combinations that do not require estimating many parameters such as arithmetic averages or weights based on the inverse mean squared forecast error – do better than more sophisticated rules relying on estimating optimal weights that depend on the full variance-covariance matrix of forecast errors, cf. Bunn (1985), Clemen and Winkler (1986), Dunis, Laws and Chauvin (2001), Figlewski and Urich (1983) and Makridakis and Winkler (1983).

Palm and Zellner (1992, p. 699) concisely summarize the advantages of adopting a simple average forecast:

- “1. Its weights are known and do not have to be estimated, an important advantage if there is little evidence on the performance of individual forecasts or if the parameters of the model generating the forecasts are time-varying;
2. In many situations a simple average of forecasts will achieve a substantial reduction in variance and bias through averaging out individual bias;

3. It will often dominate, in terms of MSE, forecasts based on optimal weighting if proper account is taken of the effect of sampling errors and model uncertainty on the estimates of the weights.”

Despite the impressive empirical track record of equal-weighted forecast combinations we stress that the theoretical justification for this method critically depends on the ratio of forecast error variances not being too far away from unity. They also depend on the correlation between forecast errors not varying too much across pairs of models. Consistent with this, [Gupta and Wilton \(1987\)](#) find that the performance of equal weighted combinations depends strongly on the relative size of the variance of the forecast errors associated with different forecasting methods. When these are similar, equal weights perform well, while when larger differences are observed, differential weighting of forecasts is generally required.

Another reason for the good average performance of equal-weighted forecast combinations is related to model instability. If model instability is sufficiently important to render precise estimation of combination weights nearly impossible, equal-weighting of forecasts may become an attractive alternative as pointed out by [Figlewski and Urich \(1983\)](#), [Clemen and Winkler \(1986\)](#), [Kang \(1986\)](#), [Diebold and Pauly \(1987\)](#) and [Palm and Zellner \(1992\)](#).

Results regarding the performance of equal-weighted forecast combinations may be sensitive to the loss function underlying the problem. [Elliott and Timmermann \(2005\)](#) find in an empirical application that the optimal weights in a combination of inflation survey forecasts and forecasts from a simple autoregressive model strongly depend on the degree of asymmetry in the loss function. In the absence of loss asymmetry, the autoregressive forecast does not add much information. However, under asymmetric loss (in either direction), both sets of forecasts appear to contain information and have non-zero weights in the combined forecast. Their application confirms the frequent finding that equal-weights outperform estimated optimal weights under MSE loss. However, it also shows very clearly that this result can be overturned under asymmetric loss where use of estimated optimal weights may lead to smaller average losses out-of-sample.

### *7.2. Choosing the single forecast with the best track record is often a bad idea*

Many studies have found that combination dominates the best individual forecast in out-of-sample forecasting experiments. For example, [Makridakis et al. \(1982\)](#) report that a simple average of six forecasting methods performed better than the underlying individual forecasts. In simulation experiments [Gupta and Wilton \(1987\)](#) also find combination superior to the single best forecast. [Makridakis and Winkler \(1983\)](#) report large gains from simply averaging forecasts from individual models over the performance of the best model. [Hendry and Clements \(2002\)](#) explain the better performance of combination methods over the best individual model by misspecification of the models caused by deterministic shifts in the underlying data generating process. Naturally, the models cannot be misspecified in the same way with regard to this source of change, or else diversification gains would be zero.



In one of the most comprehensive studies to date, [Stock and Watson \(2001\)](#) consider combinations of a range of linear and nonlinear models fitted to a very large set of US macroeconomic variables. They find strong evidence in support of using forecast combination methods, particularly the average or median forecast and the forecasts weighted by their inverse MSE. The overall dominance of the combination forecasts holds at the one, six and twelve month horizons. Furthermore, the best combination methods combine forecasts across many different time-series models.

Similarly, in a time-series simulation experiment, [Winkler and Makridakis \(1983\)](#) find that a weighted average with weights inversely proportional to the sum of squared errors or a weighted average with weights that depend on the exponentially discounted sum of squared errors perform better than the best individual forecasting model, equal-weighting or methods that require estimation of the full covariance matrix for the forecast errors.

[Aiolfi and Timmermann \(2006\)](#) find evidence of persistence in the out-of-sample performance of linear and nonlinear forecasting models fitted to a large set of macroeconomic time-series in the G7 countries. Models that were in the top and bottom quartiles when ranked by their historical forecasting performance have a higher than average chance of remaining in the top and bottom quartiles, respectively, in the out-of-sample period. They also find systematic evidence of ‘crossings’, where the previous best models become the worst models in the future or vice versa, particularly among the linear forecasting models. They find that many forecast combinations produce lower out-of-sample MSE than a strategy of selecting the previous best forecasting model irrespective of the length of the backward-looking window used to measure past forecasting performance.

### *7.3. Trimming of the worst models often improves performance*

Trimming of forecasts can occur at two levels. First, it can be adopted as a form of outlier reduction rule [cf. [Chan, Stock and Watson \(1999\)](#)] at the initial stage that produces forecasts from the individual models. Second it can be used in the combination stage where models deemed to be too poor may be discarded. Since the first form of trimming has more to do with specification of the individual models underlying the forecast combination, we concentrate on the latter form of trimming which has been used successfully in many studies. Most obviously, when many forecasts get a weight close to zero, improvements due to reduced parameter estimation errors can be gained by dropping such models.

[Winkler and Makridakis \(1983\)](#) find that including very poor models in an equal-weighted combination can substantially worsen forecasting performance. [Stock and Watson \(2004\)](#) also find that the simplest forecast combination methods such as trimmed equal weights and slowly moving weights tend to perform well and that such combinations do better than forecasts from a dynamic factor model.

In their thick modeling approach, [Granger and Jeon \(2004\)](#) recommend trimming five or ten percent of the worst models, although the extent of the trimming will depend on the application at hand.

More aggressive trimming has also been proposed. In a forecasting experiment involving the prediction of stock returns by means of a large set of forecasting models, [Aiolfi and Favero \(2005\)](#) investigate the performance of a large set of trimming schemes. Their findings indicate that the best performance is obtained when the top 20% of the forecasting models is combined in the forecast so that 80% of the models (ranked by their  $R^2$ -value) are trimmed.

#### 7.4. Shrinkage often improves performance

By and large shrinkage methods have performed quite well in empirical studies. In an empirical exercise containing four real-time forecasts of nominal and real GNP, [Diebold and Pauly \(1990\)](#) report that shrinkage weights systematically improve upon the forecasting performance over methods that select a single forecast or use least squares estimates of the combination weights. They direct the shrinkage towards a prior reflecting equal weights and find that the optimal degree of shrinkage tends to be large. Similarly, [Stock and Watson \(2004\)](#) find that shrinkage methods perform best when the degree of shrinkage (towards equal weights) is quite strong.

[Aiolfi and Timmermann \(2006\)](#) explore persistence in the performance of forecasting models by proposing a set of combination strategies that first pre-select models into either quartiles or clusters on the basis of the distribution of past forecasting performance across models. Then they pool forecasts within each cluster and estimate optimal combination weights that are shrunk towards equal weights. These conditional combination strategies lead to better average forecasting performance than simpler strategies in common use such as using the single best model or averaging across all forecasting models or a small subset of these.

[Elliott \(2004\)](#) undertakes a simulation experiment where he finds that although shrinkage methods always dominate least squares estimates of the combination weights, the performance of the shrinkage method can be quite sensitive to the shrinkage parameter and that none of the standard methods for determining this parameter work particularly well.

Given the similarity of the mean-variance optimization problem in finance to the forecast combination problem, it is not surprising that empirical findings in finance mirror those in the forecast combination literature. For example, it has generally been found in applications to asset returns that sample estimates of portfolio weights that solve a standard mean-variance optimization problem are extremely sensitive to small changes in sample means. In addition they are highly sensitive to variations in the inverse of the covariance matrix estimate,  $\hat{\Sigma}^{-1}$ .

[Jobson and Korkie \(1980\)](#) show that the sample estimate of the optimal portfolio weights can be characterized as the ratio of two estimators, each of whose first and second moments can be derived in closed form. They use Taylor series expansions to

derive an approximate solution for the first two moments of the optimal weights, noting that higher order moments can be characterized under additional normality assumptions. They also derive the asymptotic distribution of the portfolio weights for the case where  $N$  is fixed and  $T$  goes to infinity. In simulation experiments they demonstrate that the sample estimates of the portfolio weights are highly volatile and can take extreme values that lead to poor out-of-sample performance.

It is widely recognized in finance that imposing portfolio weight constraints generally leads to improved out-of-sample performance of mean-variance efficient portfolios. For example, [Jaganathan and Ma \(2003\)](#) find empirically that once such constraints are imposed on portfolio weights, other refinements of covariance matrix estimation have little additional effect on the variance of the optimal portfolio. Since they also demonstrate that portfolio weight constraints can be interpreted as a form of shrinkage, these findings lend support to using shrinkage methods as well.

Similarly, [Ledoit and Wolf \(2003\)](#) report that the out-of-sample standard deviation of portfolio returns based on a shrunk covariance matrix is significantly lower than the standard deviation of portfolio returns based on more conventional estimates of the covariance matrix.

Notice that shrinkage and trimming tend to work in opposite directions – at least if the shrinkage is towards equal weights. Shrinkage tends to give more similar weights to all models whereas trimming completely discards a subset of models. If some models produce extremely poor out-of-sample forecasts, shrinkage can be expected to perform poorly if the combined forecast is shrunk too aggressively towards an equal-weighted average. For this reason, shrinkage preceded by a trimming step may work well in many situations.

### *7.5. Limited time-variation in the combination weights may be helpful*

Empirical evidence on the value of allowing for time-varying combinations in the combination weights is somewhat mixed. Time-variations in forecasts can be introduced either in the individual models underlying the combination or in the combination weights themselves and both approaches have been considered. The idea of time-varying forecast combinations goes back to the advent of the combination literature in economics. [Bates and Granger \(1969\)](#) used combination weights that were adaptively updated as did many subsequent studies such as [Winkler and Makridakis \(1983\)](#). [Newbold and Granger \(1974\)](#) considered values of the window length,  $v$ , in (47) and (48) between one and twelve periods and values of the discounting factor,  $\lambda$ , in (50) and (51) between 1 and 2.5. Their results suggested that there is an interior optimum around  $v = 6$ ,  $\alpha = 0.5$  for which the adaptive updating method (49) performs best whereas the rolling window combinations generally do best for the longest windows, i.e.,  $v = 9$  or  $v = 12$ , and the best exponential discounting was found for  $\lambda$  around 2 or 2.5. This is consistent with the finding by [Bates and Granger \(1969\)](#) that high values of the discounting factor tend to work best. A method that combines a Holt–Winters and stepwise autoregressive forecast was found to perform particularly well. [Winkler and Makridakis \(1983\)](#) report similar

results and also find that the longer windows,  $v$ , in equations such as (47) and (48) tend to produce the most accurate forecasts, although in their study the best results among the discounting methods were found for relatively low values of the discount factor.

In a combination of forecasts from the Survey of Professional Forecasters and forecasts from simple autoregressive models applied to six macroeconomic variables, Elliott and Timmermann (2005) investigate the out-of-sample forecasting performance produced by different constant and time-varying forecasting schemes such as (57). Compared to a range of other time-varying forecast combination methods, a two-state regime switching method produces a lower MSE-value for four or five out of six cases. They argue that the evidence suggests that the best forecast combination method allows the combination weights to vary over time but in a mean-reverting manner. Unsurprisingly, allowing for three states leads to worse forecasting performance for four of the six variables under consideration.

Stock and Watson (2004) report that the combined forecasts that perform best in their study are the time-varying parameter (TVP) forecast with very little time variation, the simple mean and a trimmed mean. They conclude that “the results for the methods designed to handle time variation are mixed. The TVP forecasts sometimes work well but sometimes work quite poorly and in this sense are not robust; the larger the amount of time variation, the less robust are the forecasts. Similarly, the discounted MSE forecasts with the most discounting ... are typically no better than, and sometimes worse than, their counterparts with less or no discounting.”

This leads them to conclude that “This “forecast combination puzzle” – the repeated finding that simple combination forecasts outperform sophisticated adaptive combination methods in empirical applications – is, we think, more likely to be understood in the context of a model in which there is widespread instability in the performance of individual forecast, but the instability is sufficiently idiosyncratic that the combination of these individually unstably performing forecasts can itself be stable.”

### 7.6. Empirical application

To demonstrate the practical use of forecast combination techniques, we consider an empirical application to the seven-country data set introduced in Stock and Watson (2004). This data comprises up to 43 quarterly time series for each of the G7 economies (Canada, France, Germany, Italy, Japan, UK, and the US) over the period 1959Q1–1999Q4. Observations on some variables are only available for a shorter sample. The 43 series comprise the following categories: Asset returns, interest rates and spreads; measures of real economic activity; prices and wages; and various monetary aggregates. The data has been transformed as described in Stock and Watson (2004) and Aiolfi and Timmermann (2006) to deal with seasonality, outliers and stochastic trends, yielding between 46 and 71 series per country.

Forecasts are generated from bivariate autoregressive models of the type

$$y_{t+h} = c + A(L)y_t + B(L)x_t + \epsilon_{t+h}, \quad (82)$$

where  $x_t$  is a regressor other than  $y_t$ . Lag lengths are selected recursively using the BIC with between 1 and 4 lags of  $x_t$  and between 0 and 4 lags of  $y_t$ . All parameters are estimated recursively using an expanding data window. For more details, see Aiolfi and Timmermann (2006). The average number of forecasting models entertained ranges from 36 for France, through 67 for the US.

We consider three trimmed forecast combination schemes that take simple averages over the top 25%, top 50% and top 75% of forecast models ranked recursively by means of their forecasting performance up to the point in time where a new out-of-sample forecast gets computed. In addition we report the performance of the simple average (mean) forecast, the median forecast, the triangular forecast combination scheme (38) and the discounted mean squared forecast combination (50) with  $\lambda = 1$  so the forecasting models get weighted by the inverse of their MSE-values. Out-of-sample forecasting performance is reported relative to the forecasting performance of the previous best (PB) model selected according to the forecasting performance up to the point where a new out-of-sample forecast is generated. Numbers below one indicate better MSE performance while numbers above one indicate worse performance relative to this benchmark. The out-of-sample period is 1970Q1–1999Q4.

Table 2 reports results averaged across variables within each country.<sup>15</sup> We show results for four forecast horizons, namely  $h = 1, 2, 4$  and 8. For each country it is clear that simple trimmed forecast combinations perform very well and generally are better, the fewer models that get included, i.e. the more aggressive the trimming. Furthermore, gains can be quite large – on the order of 10–15% relative to the forecast from the previous best model. The median forecast also performs better on average than the previous best model, but is generally worse compared to some of the other combination schemes as is the discounted mean squared forecast error weighting scheme. Results are quite consistent across the seven countries.

Table 3 shows results averaged across countries but for the separate categories of variables. Gains from forecast combination tend to be greater for the economic activity variables and somewhat smaller for the monetary aggregates. There is also a systematic tendency for the forecasting performance of the combinations relative to the best single model to improve as the forecast horizon is extended from one-quarter to two or more quarters.

How consistent are these results across countries and variables? To investigate this question, Tables 4, 5 and 6 show disaggregate results for the US, Japan and France. Considerable variations in gains from forecast combinations emerge across countries, variables and horizons. Table 4 shows that gains in the US are very large for the economic activity variables but somewhat smaller for asset returns, interest rates and monetary aggregates. Compared to the US results, in Japan the best combinations perform relatively worse for economic activity variables and prices and wages but relatively better for the monetary aggregates, asset returns and interest rates. Finally in the case of

<sup>15</sup> I am grateful to Marco Aiolfi for carrying out these calculations.

Table 2

*Linear Models.* Out-of-sample forecasting performance of combination schemes applied to linear models. Each panel reports the out-of-sample MSFE – relative to that of the previous best (PB) model using an expanding window – averaged across variables, for different combination strategies, countries and forecast horizons ( $h$ ).

$h = 1$								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
US	0.88	0.89	0.90	0.90	0.93	0.90	0.91	1.00
UK	0.91	0.91	0.92	0.92	0.93	0.91	0.92	1.00
Germany	0.92	0.93	0.93	0.92	0.95	0.92	0.92	1.00
Japan	0.93	0.94	0.94	0.94	0.97	0.94	0.94	1.00
Italy	0.90	0.90	0.91	0.91	0.93	0.90	0.91	1.00
France	0.93	0.93	0.94	0.94	0.96	0.93	0.94	1.00
Canada	0.91	0.91	0.92	0.92	0.94	0.91	0.92	1.00
$h = 2$								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
US	0.85	0.86	0.86	0.86	0.88	0.86	0.86	1.00
UK	0.90	0.90	0.90	0.91	0.92	0.90	0.91	1.00
Germany	0.90	0.90	0.91	0.91	0.93	0.90	0.91	1.00
Japan	0.90	0.91	0.92	0.92	0.94	0.91	0.92	1.00
Italy	0.89	0.89	0.89	0.89	0.90	0.89	0.89	1.00
France	0.88	0.88	0.88	0.88	0.89	0.88	0.88	1.00
Canada	0.90	0.90	0.91	0.90	0.94	0.90	0.90	1.00
$h = 4$								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
US	0.87	0.87	0.87	0.87	0.90	0.87	0.87	1.00
UK	0.86	0.86	0.86	0.86	0.87	0.86	0.86	1.00
Germany	0.90	0.90	0.91	0.91	0.92	0.90	0.91	1.00
Japan	0.91	0.93	0.95	0.96	0.98	0.94	0.97	1.00
Italy	0.86	0.85	0.85	0.85	0.86	0.85	0.85	1.00
France	0.88	0.88	0.88	0.88	0.89	0.88	0.88	1.00
Canada	0.85	0.85	0.86	0.86	0.88	0.85	0.86	1.00
$h = 8$								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
US	0.85	0.85	0.86	0.86	0.88	0.85	0.86	1.00
UK	0.88	0.88	0.89	0.89	0.91	0.88	0.89	1.00
Germany	0.90	0.91	0.91	0.91	0.92	0.90	0.91	1.00
Japan	0.85	0.85	0.85	0.85	0.86	0.85	0.85	1.00
Italy	0.89	0.89	0.90	0.90	0.91	0.89	0.90	1.00
France	0.90	0.90	0.90	0.90	0.92	0.90	0.90	1.00
Canada	0.86	0.87	0.87	0.87	0.88	0.86	0.86	1.00

Note: TMB25%, TMB50% and TMB75% use the mean forecast computed across the top 25%, 50% and 75% of models ranked by historical forecasting performance. Mean and median use the mean or median forecast across all models. TK is the forecast from a triangular weighting scheme (38), while DMSFE is the forecast produced by the discounted mean squared forecast error scheme in (50).

Table 3

*Linear Models.* Out-of-sample forecasting performance of combination schemes applied to linear models. Each panel reports the out-of-sample MSFE – relative to that of the previous best model using an expanding window – averaged across countries, for different combination strategies, categories of economic variables and forecast horizons ( $h$ ).

	All							
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.91	0.92	0.92	0.92	0.94	0.92	0.92	1.00
$h = 2$	0.89	0.89	0.89	0.89	0.91	0.89	0.90	1.00
$h = 4$	0.88	0.88	0.88	0.88	0.90	0.88	0.89	1.00
$h = 8$	0.87	0.88	0.88	0.88	0.90	0.88	0.88	1.00
Returns and interest rates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.92	0.92	0.92	0.92	0.94	0.92	0.92	1.00
$h = 2$	0.89	0.90	0.90	0.90	0.91	0.90	0.90	1.00
$h = 4$	0.88	0.89	0.89	0.89	0.91	0.88	0.89	1.00
$h = 8$	0.87	0.87	0.87	0.87	0.89	0.87	0.87	1.00
Economic activity								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.89	0.91	0.92	0.93	0.95	0.91	0.93	1.00
$h = 2$	0.86	0.88	0.89	0.89	0.93	0.88	0.90	1.00
$h = 4$	0.85	0.88	0.89	0.89	0.93	0.88	0.90	1.00
$h = 8$	0.87	0.89	0.90	0.91	0.95	0.89	0.90	1.00
Prices and wages								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.90	0.91	0.91	0.91	0.93	0.91	0.91	1.00
$h = 2$	0.89	0.89	0.89	0.89	0.91	0.89	0.89	1.00
$h = 4$	0.86	0.86	0.87	0.87	0.88	0.86	0.87	1.00
$h = 8$	0.87	0.86	0.86	0.86	0.88	0.86	0.86	1.00
Monetary aggregates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.91	0.92	0.93	0.93	0.96	0.92	0.93	1.00
$h = 2$	0.89	0.89	0.89	0.89	0.90	0.89	0.89	1.00
$h = 4$	0.90	0.90	0.90	0.89	0.90	0.89	0.89	1.00
$h = 8$	0.90	0.90	0.90	0.90	0.91	0.90	0.90	1.00

Note: see note of Table 2.

Table 4

*Linear models US.* Out-of-sample forecasting performance of combination schemes applied to linear models. Each panel reports the out-of-sample MSFE – relative to that of the previous best model using an expanding window – averaged across variables, for different combination strategies, categories of economic variables and forecast horizons ( $h$ ).

	All							
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.88	0.89	0.90	0.90	0.93	0.90	0.91	1.00
$h = 2$	0.85	0.86	0.86	0.86	0.88	0.86	0.86	1.00
$h = 4$	0.87	0.87	0.87	0.87	0.90	0.87	0.87	1.00
$h = 8$	0.85	0.85	0.86	0.86	0.88	0.85	0.86	1.00
Returns and interest rates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.89	0.89	0.89	0.89	0.91	0.89	0.89	1.00
$h = 2$	0.87	0.87	0.88	0.88	0.90	0.87	0.88	1.00
$h = 4$	0.90	0.90	0.90	0.90	0.92	0.90	0.90	1.00
$h = 8$	0.86	0.86	0.86	0.86	0.87	0.86	0.86	1.00
Economic activity								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.86	0.90	0.91	0.92	0.94	0.90	0.92	1.00
$h = 2$	0.77	0.80	0.81	0.82	0.87	0.80	0.82	1.00
$h = 4$	0.80	0.83	0.84	0.84	0.90	0.83	0.84	1.00
$h = 8$	0.82	0.86	0.88	0.90	0.98	0.86	0.88	1.00
Prices and wages								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.86	0.86	0.87	0.87	0.90	0.86	0.87	1.00
$h = 2$	0.84	0.85	0.84	0.85	0.86	0.84	0.85	1.00
$h = 4$	0.83	0.83	0.83	0.82	0.83	0.83	0.82	1.00
$h = 8$	0.80	0.79	0.79	0.79	0.81	0.79	0.79	1.00
Monetary aggregates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.92	0.95	0.97	0.98	1.03	0.96	0.98	1.00
$h = 2$	0.88	0.88	0.87	0.87	0.88	0.87	0.88	1.00
$h = 4$	0.87	0.88	0.88	0.88	0.90	0.88	0.88	1.00
$h = 8$	0.93	0.92	0.93	0.93	0.94	0.92	0.93	1.00

Note: see note of Table 2.



Table 5

*Linear models: Japan.* Out-of-sample forecasting performance of combination schemes applied to linear models. Each panel reports the out-of-sample MSFE – relative to that of the previous best model using an expanding window – averaged across variables, for different combination strategies, categories of economic variables and forecast horizons ( $h$ ).

	All							
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.93	0.94	0.94	0.94	0.97	0.94	0.94	1.00
$h = 2$	0.90	0.91	0.92	0.92	0.94	0.91	0.92	1.00
$h = 4$	0.91	0.93	0.95	0.96	0.98	0.94	0.97	1.00
$h = 8$	0.85	0.85	0.85	0.85	0.86	0.85	0.85	1.00
Returns and interest rates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.94	0.95	0.96	0.96	1.00	0.95	0.96	1.00
$h = 2$	0.92	0.93	0.93	0.93	0.95	0.93	0.94	1.00
$h = 4$	0.91	0.93	0.94	0.95	0.98	0.93	0.96	1.00
$h = 8$	0.81	0.81	0.82	0.82	0.83	0.81	0.82	1.00
Economic activity								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.97	0.99	1.00	1.00	1.02	0.99	1.00	1.00
$h = 2$	0.91	0.93	0.94	0.95	0.96	0.93	0.95	1.00
$h = 4$	0.99	1.00	1.03	1.05	1.06	1.01	1.06	1.00
$h = 8$	0.89	0.88	0.88	0.89	0.89	0.88	0.88	1.00
Prices and wages								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.90	0.92	0.93	0.92	0.94	0.92	0.92	1.00
$h = 2$	0.91	0.93	0.93	0.93	0.97	0.92	0.93	1.00
$h = 4$	0.90	0.95	0.98	0.99	1.03	0.96	1.00	1.00
$h = 8$	0.90	0.90	0.89	0.89	0.91	0.89	0.90	1.00
Monetary aggregates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.89	0.90	0.89	0.89	0.91	0.89	0.89	1.00
$h = 2$	0.85	0.85	0.85	0.85	0.86	0.85	0.85	1.00
$h = 4$	0.87	0.87	0.87	0.87	0.88	0.87	0.86	1.00
$h = 8$	0.84	0.83	0.83	0.83	0.83	0.83	0.83	1.00

Note: see note of Table 2.

Table 6

*Linear models: France.* Out-of-sample forecasting performance of combination schemes applied to linear models. Each panel reports the out-of-sample MSFE – relative to that of the previous best model using an expanding window – averaged across variables, for different combination strategies, categories of economic variables and forecast horizons ( $h$ ).

All								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.93	0.93	0.94	0.94	0.96	0.93	0.94	1.00
$h = 2$	0.88	0.88	0.88	0.88	0.89	0.88	0.88	1.00
$h = 4$	0.88	0.88	0.88	0.88	0.89	0.88	0.88	1.00
$h = 8$	0.90	0.90	0.90	0.90	0.92	0.90	0.90	1.00
Returns and interest rates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.94	0.94	0.95	0.94	0.97	0.94	0.95	1.00
$h = 2$	0.89	0.89	0.89	0.89	0.89	0.89	0.89	1.00
$h = 4$	0.89	0.89	0.89	0.89	0.90	0.89	0.89	1.00
$h = 8$	0.89	0.89	0.90	0.89	0.91	0.89	0.90	1.00
Economic activity								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.80	0.80	0.81	0.82	0.85	0.80	0.83	1.00
$h = 2$	0.75	0.76	0.77	0.77	0.79	0.76	0.77	1.00
$h = 4$	0.78	0.77	0.77	0.78	0.78	0.77	0.77	1.00
$h = 8$	0.84	0.84	0.84	0.84	0.86	0.83	0.84	1.00
Prices and wages								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.96	0.96	0.96	0.97	0.98	0.96	0.97	1.00
$h = 2$	0.90	0.90	0.91	0.90	0.92	0.90	0.90	1.00
$h = 4$	0.86	0.85	0.85	0.85	0.86	0.85	0.85	1.00
$h = 8$	0.91	0.90	0.90	0.91	0.93	0.90	0.91	1.00
Monetary aggregates								
	TMB25%	TMB50%	TMB75%	Mean	Median	TK	DMSFE	PB
$h = 1$	0.88	0.89	0.91	0.91	0.94	0.90	0.91	1.00
$h = 2$	0.85	0.86	0.86	0.87	0.90	0.86	0.87	1.00
$h = 4$	1.06	1.07	1.08	1.09	1.11	1.07	1.09	1.00
$h = 8$	0.99	1.01	1.01	1.01	1.05	1.00	1.01	1.00

Note: see note of Table 2.

France, we uncover a number of cases where, for the forecasts of monetary aggregates, in fact none of the combinations beat the previous best model.

## 8. Conclusion

In his classical survey of forecast combinations, Clemen (1989, p. 567) concluded that “Combining forecasts has been shown to be practical, economical and useful. Underlying theory has been developed, and many empirical tests have demonstrated the value of composite forecasting. We no longer need to justify this methodology.”

In the early days of the combination literature the set of forecasts was often taken as given, but recent experiments undertaken by Stock and Watson (2001, 2004) and Marcellino (2004) let the forecast user control both the number of forecasting models as well as the types of forecasts that are being combined. This opens a whole new set of issues: is it best to combine forecasts from linear models with different regressors or is it better to combine forecasts produced by different families of models, e.g., linear and nonlinear, or maybe the same model using estimators with varying degrees of robustness? The answer to this depends of course on the type of misspecification or instability the model combination can hedge against. Unfortunately this is typically unknown so general answers are hard to come by.

Since then, combination methods have gained even more ground in the forecasting literature, largely because of the strength of the empirical evidence suggesting that these methods systematically perform better than alternatives based on forecasts from a single model. Stable, equal weights have so far been the workhorse of the combination literature and have set a benchmark that has proved surprisingly difficult to beat. This is surprising since – on theoretical grounds – one would not expect any particular combination scheme to be dominant, since the various methods incorporate restrictions on the covariance matrix that are designed to trade off bias against reduced parameter estimation error. The optimal bias can be expected to vary across applications, and the scheme that provides the best trade-off is expected to depend on the sample size, the number of forecasting models involved, the ratio of the variance of individual models’ forecast errors as well as their correlations and the degree of instability in the underlying data generating process.

Current research also provides encouraging pointers towards modifications of this simple strategy that can improve forecasting. Modest time-variations in the combination weights and trimming of the worst models have generally been found to work well, as has shrinkage towards equal weights or some other target requiring the estimation of a relatively modest number of parameters, particularly in applications with combinations of a large set of forecasts.

## Acknowledgements

This research was sponsored by NSF grant SES0111238. I am grateful to Marco Aiolfi, Graham Elliott and Clive Granger for many discussions on the topic. Barbara Rossi

and Mark Watson provided detailed comments and suggestions that greatly improved the paper. Comments from seminar participants at the UCSD Rady School forecasting conference were also helpful.

## References

- Aiolfi, M., Favero, C.A. (2005). "Model uncertainty, thick modeling and the predictability of stock returns". *Journal of Forecasting* 24, 233–254.
- Aiolfi, M., Timmermann, A. (2006). "Persistence of forecasting performance and combination strategies". *Journal of Econometrics*. In press.
- Armstrong, J.S. (1989). "Combining forecasts: The end of the beginning or the beginning of the end". *International Journal of Forecasting* 5, 585–588.
- Bates, J.M., Granger, C.W.J. (1969). "The combination of forecasts". *Operations Research Quarterly* 20, 451–468.
- Bunn, D.W. (1975). "A Bayesian approach to the linear combination of forecasts". *Operations Research Quarterly* 26, 325–329.
- Bunn, D.W. (1985). "Statistical efficiency in the linear combination of forecasts". *International Journal of Forecasting* 1, 151–163.
- Chan, Y.L., Stock, J.H., Watson, M.W. (1999). "A dynamic factor model framework for forecast combination". *Spanish Economic Review* 1, 91–122.
- Chong, Y.Y., Hendry, D.F. (1986). "Econometric evaluation of linear macro-economic models". *Review of Economic Studies* 53, 671–690.
- Christoffersen, P., Diebold, F.X. (1997). "Optimal prediction under asymmetrical loss". *Econometric Theory* 13, 806–817.
- Clemen, R.T. (1987). "Combining overlapping information". *Management Science* 33, 373–380.
- Clemen, R.T. (1989). "Combining forecasts: A review and annotated bibliography". *International Journal of Forecasting* 5, 559–581.
- Clemen, R.T., Murphy, A.H., Winkler, R.L. (1995). "Screening probability forecasts: Contrasts between choosing and combining". *International Journal of Forecasting* 11, 133–145.
- Clemen, R.T., Winkler, R.L. (1986). "Combining economic forecasts". *Journal of Business and Economic Statistics* 4, 39–46.
- Deutsch, M., Granger, C.W.J., Terasvirta, T. (1994). "The combination of forecasts using changing weights". *International Journal of Forecasting* 10, 47–57.
- Diebold, F.X. (1988). "Serial correlation and the combination of forecasts". *Journal of Business and Economic Statistics* 6, 105–111.
- Diebold, F.X. (1989). "Forecast combination and encompassing: Reconciling two divergent literatures". *International Journal of Forecasting* 5, 589–592.
- Diebold, F.X., Lopez, J.A. (1996). "Forecast evaluation and combination". In: Maddala, G.S., Rao, C.R. (Eds.), *Statistical Methods in Finance, Handbook of Statistics*, vol. 14. Elsevier, Amsterdam, pp. 241–268.
- Diebold, F.X., Pauly, P. (1987). "Structural change and the combination of forecasts". *Journal of Forecasting* 6, 21–40.
- Diebold, F.X., Pauly, P. (1990). "The use of prior information in forecast combination". *International Journal of Forecasting* 6, 503–508.
- Donaldson, R.G., Kamstra, M. (1996). "Forecast combining with neural networks". *Journal of Forecasting* 15, 49–61.
- Dunis, C., Laws, J., Chauvin, S. (2001). "The use of market data and model combinations to improve forecast accuracy". In: Dunis, C., Timmermann, A., Moody, J.E. (Eds.), *Development in Forecasts Combination and Portfolio Choice*. Wiley, Oxford.

- Dunis, C.L., Timmermann, A., Moody, J.E. (Eds.) (2001). *Developments in Forecasts Combination and Portfolio Choice*. Wiley, Oxford.
- Elliott, G. (2004). "Forecast combination with many forecasts". Mimeo, Department of Economics, University of California, San Diego.
- Elliott, G., Timmermann, A. (2004). "Optimal forecast combinations under general loss functions and forecast error distributions". *Journal of Econometrics* 122, 47–79.
- Elliott, G., Timmermann, A. (2005). "Optimal forecast combination weights under regime switching". *International Economic Review* 46, 1081–1102.
- Engle, R.F., Granger, C.W.J., Kraft, D. (1984). "Combining competing forecasts of inflation using a bivariate ARCH model". *Journal of Economic Dynamics and Control* 8, 151–165.
- Figlewski, S., Urlich, T. (1983). "Optimal aggregation of money supply forecasts: Accuracy, profitability and market efficiency". *Journal of Finance* 28, 695–710.
- Genest, S., Zidek, J. (1986). "Combining probability distributions: A critique and an annotated bibliography". *Statistical Science* 1, 114–148.
- Geweke, J., Whitmann, C. (2006). "Bayesian forecasting". In: Elliott, G., Granger, C.W.J., Timmermann, A. (Eds.), *Handbook of Economic Forecasting*. Elsevier, Amsterdam, pp. 3–79. Chapter 1 in this volume.
- Giacomini, R., Komunjer, I. (2005). "Evaluation and combination of conditional quantile forecasts". *Journal of Business and Economic Statistics* 23, 416–431.
- Granger, C.W.J., Jeon, Y. (2004). "Thick modeling". *Economic Modelling* 21, 323–343.
- Granger, C.W.J., Machina, M.J. (2006). "Forecasting and decision theory". In: Elliott, G., Granger, C.W.J., Timmermann, A. (Eds.), *Handbook of Economic Forecasting*. Elsevier, Amsterdam, pp. 81–98. Chapter 2 in this volume.
- Granger, C.W.J., Pesaran, M.H. (2000). "Economic and statistical measures of forecast accuracy". *Journal of Forecasting* 19, 537–560.
- Granger, C.W.J., Ramanathan, R. (1984). "Improved methods of combining forecasts". *Journal of Forecasting* 3, 197–204.
- Guidolin, M., Timmermann, A. (2005). "Optimal forecast combination weights under regime shifts with an application to US interest rates". Mimeo, Federal Reserve Bank of St. Louis and the Department of Economics, University of California, San Diego.
- Gupta, S., Wilton, P.C. (1987). "Combination of forecasts: An extension". *Management Science* 33, 356–372.
- Hamilton, J.H. (1989). "A new approach to the economic analysis of nonstationary time series and the business cycle". *Econometrica* 57, 357–384.
- Hendry, D.F., Clements, M.P. (2002). "Pooling of forecasts". *Econometrics Journal* 5, 1–26.
- Hoeting, J.A., Madigan, D., Raftery, A.E., Volinsky, C.T. (1999). "Bayesian model averaging: A tutorial". *Statistical Science* 14, 382–417.
- Jackson, T., Karlsson, S. (2004). "Finding good predictors for inflation: A Bayesian model averaging approach". *Journal of Forecasting* 23, 479–498.
- Jaganathan, R., Ma, T. (2003). "Risk reduction in large portfolios: Why imposing the wrong constraints helps". *Journal of Finance* 58, 1651–1684.
- Jobson, J.D., Korkie, B. (1980). "Estimation for Markowitz efficient portfolios". *Journal of the American Statistical Association* 75, 544–554.
- Kang, H. (1986). "Unstable weights in the combination of forecasts". *Management Science* 32, 683–695.
- Leamer, E. (1978). *Specification Searches*. Wiley, Oxford.
- Ledoit, O., Wolf, M. (2003). "Improved estimation of the covariance matrix of stock returns with an application to portfolio election". *Journal of Empirical Finance* 10, 603–621.
- Ledoit, O., Wolf, M. (2004). "Honey, I shrunk the sample covariance matrix". *Journal of Portfolio Management*.
- LeSage, J.P., Magura, M. (1992). "A mixture-model approach to combining forecasts". *Journal of Business and Economic Statistics* 10, 445–453.
- Makridakis, S. (1989). "Why combining works?". *International Journal of Forecasting* 5, 601–603.
- Makridakis, S., Hibon, M. (2000). "The M3-competition: Results, conclusions and implications". *International Journal of Forecasting* 16, 451–476.

- Makridakis, S., Winkler, R.L. (1983). "Averages of forecasts: Some empirical results". *Management Science* 29, 987–996.
- Makridakis, S., Andersen, A., Carbone, R., Fildes, R., Hibon, M., Lewandowski, R., Newton, J., Parzen, E., Winkler, R. (1982). "The accuracy of extrapolation (time series) methods: Results of a forecasting competition". *Journal of Forecasting* 1, 111–153.
- Marcellino, M. (2004). "Forecast pooling for short time series of macroeconomic variables". *Oxford Bulletin of Economic and Statistics* 66, 91–112.
- Min, C.-K., Zellner, A. (1993). "Bayesian and non-Bayesian methods for combining models and forecasts with applications to forecasting international growth rates". *Journal of Econometrics* 56, 89–118.
- Newbold, P., Granger, C.W.J. (1974). "Experience with forecasting univariate time series and the combination of forecasts". *Journal of the Royal Statistical Society Series A* 137, 131–146.
- Newbold, P., Harvey, D.I. (2001). "Forecast combination and encompassing". In: Clements, M.P., Hendry, D.F. (Eds.), *A Companion to Economic Forecasting*. Blackwell, Oxford.
- Palm, F.C., Zellner, A. (1992). "To combine or not to combine? Issues of combining forecasts". *Journal of Forecasting* 11, 687–701.
- Patton, A., Timmermann, A. (2004). "Properties of optimal forecasts under asymmetric loss and nonlinearity". Mimeo, London School of Economics and Department of Economics, University of California, San Diego.
- Pesaran, M.H., Timmermann, A. (2005). "Selection of estimation window in the presence of breaks". Mimeo, Cambridge University and Department of Economics, University of California, San Diego.
- Raftery, A.E., Madigan, D., Hoeting, J.A. (1997). "Bayesian model averaging for linear regression models". *Journal of the American Statistical Association* 92, 179–191.
- Reid, D.J. (1968). "Combining three estimates of gross domestic product". *Economica* 35, 431–444.
- Sanders, F. (1963). "On subjective probability forecasting". *Journal of Applied Meteorology* 2, 196–201.
- Sessions, D.N., Chattererjee, S. (1989). "The combining of forecasts using recursive techniques with nonstationary weights". *Journal of Forecasting* 8, 239–251.
- Stock, J.H., Watson, M. (2001). "A comparison of linear and nonlinear univariate models for forecasting macroeconomic time series". In: Engle, R.F., White, H. (Eds.), *Festschrift in Honour of Clive Granger*. Cambridge University Press, Cambridge, pp. 1–44.
- Stock, J.H., Watson, M. (2004). "Combination forecasts of output growth in a seven-country data set". *Journal of Forecasting* 23, 405–430.
- Swanson, N.R., Zeng, T. (2001). "Choosing among competing econometric forecasts: Regression-based forecast combination using model selection". *Journal of Forecasting* 6, 425–440.
- West, K.D. (2006). "Forecast evaluation". In: Elliott, G., Granger, C.W.J., Timmermann, A. (Eds.), *Handbook of Economic Forecasting*. Elsevier, Amsterdam, pp. 99–134. Chapter 3 in this volume.
- White, H. (2006). "Approximate nonlinear forecasting methods". In: Elliott, G., Granger, C.W.J., Timmermann, A. (Eds.), *Handbook of Economic Forecasting*. Elsevier, Amsterdam, pp. 459–512. Chapter 9 in this volume.
- Winkler, R.L. (1981). "Combining probability distributions from dependent information sources". *Management Science* 27, 479–488.
- Winkler, R.L. (1989). "Combining forecasts: A philosophical basis and some current issues". *International Journal of Forecasting* 5, 605–609.
- Winkler, R.L., Makridakis, S. (1983). "The combination of forecasts". *Journal of the Royal Statistical Society Series A* 146, 150–157.
- Wright, S.M., Satchell, S.E. (2003). "Generalized mean-variance analysis and robust portfolio diversification". In: Satchell, S.E., Scowcroft, A. (Eds.), *Advances in Portfolio Construction and Implementation*. Butterworth Heinemann, London, pp. 40–54.
- Yang, Y. (2004). "Combining forecasts procedures: Some theoretical results". *Econometric Theory* 20, 176–190.
- Zellner, A. (1986). "Bayesian estimation and prediction using asymmetric loss functions". *Journal of the American Statistical Association* 81, 446–451.
- Zellner, A., Hong, C., Min, C.-k. (1991). "Forecasting turning points in international output growth rates using Bayesian exponentially weighted autoregression, time-varying parameter, and pooling techniques". *Journal of Econometrics* 49, 275–304.